

STEENROD OPERATIONS AND HOCHSCHILD HOMOLOGY.

by

BITJONG NDOMBOL AND JEAN-CLAUDE THOMAS

University of DSCHANG - Cameroun and University of ANGERS - CNRS-6093
Département de Mathématiques - 2, Bd Lavoisier 49045 Angers - France.

Abstract. Let X be a simply connected space and $\mathbb{F}_p$ be a prime field. The algebra of normalized singular cochains $N^*(X; \mathbb{F}_p)$ admits a natural homotopy structure which induces natural Steenrod operations on the Hochschild homology $HH_*N^*(X; \mathbb{F}_p)$ of the space X . The primary purpose of this paper is to prove that the J. Jones isomorphism $HH_*N^*(X; \mathbb{F}_p) \cong H^*(X^{S^1}; \mathbb{F}_p)$ identifies these Steenrod operations with those defined on the cohomology of the free loop space with coefficients in $\mathbb{F}_p$. The other goal of this paper is to describe a theoretic model which allows to do some computations.

AMS Classification: 55P35, 13D03, 55P48, 16E45, 18F25.

Keywords: Hochschild homology, free loop space, Steenrod operations, minimal model.

Introduction.

Let $A = \{A^i\}_{i \geq 0}$ be a augmented differential graded algebra over the field $\mathbb{F}_p$. The homology of the normalized Hochschild **chain** complex $\mathfrak{C}_*A$ is called the Hochschild homology (with coefficients in A) of (A, d_A) and is denoted by HH_*A . Let X be a simply connected space. In 1987, J.D.S. Jones [10] constructed an isomorphism of graded vector spaces $HH_*N^*X \cong H^*(X^{S^1}; \mathbb{F}_p)$, where X^{S^1} denotes the free loop space and N^*X denotes singular cochains with coefficients in $\mathbb{F}_p$. In view of this result, one may ask the following question:

Does there exist a subcategory $\mathbf{DA}'$ of the category $\mathbf{DA}$ of differential graded algebras such that:

- (Q.1) *for any connected space X , N^*X is an object of $\mathbf{DA}'$,*
- (Q.2) *if A is an object of $\mathbf{DA}'$ then HH_*A is an $\mathcal{B}_p$ -unstable algebra,*
- (Q.3) *the Jones' isomorphism is an homomorphism of $\mathcal{A}_p$ -unstable algebras?*

Here $\mathcal{B}_p$ denotes the large mod p -Steenrod algebra and $\mathcal{A}_p = \mathcal{B}_p/(P^0 = id)$ denotes the usual mod p -Steenrod algebra.

In this paper we give an affirmative answer to (Q.1) and (Q.3) and a partial answer to (Q.2). For this purpose we introduce the notion of π -shc algebra where π denotes the cyclic group of order a fixed prime p (1.6). More precisely we prove:

Theorem A. *Let p be a prime and A be the mod p reduction of a graded module over $\mathbb{Z}$ and denote by β the Bockstein homomorphism. If A is a π -shc algebra over $\mathbb{F}_p$ then for any $i \in \mathbb{Z}$ and any $x \in (HH_*A)^n$ there exist well defined homology classes $Sq^i(x) \in (HH_*A)^{n+i}$*

if $p = 2$ and $P^i(x) \in (HH_*A)^{n+2i(p-1)}$ if $p > 2$ such that: $Sq^1(x) = P^i(1_{HA}) = 0, i \neq 0$ and $Sq^i(x) = \begin{cases} 0 & \text{if } i > n \\ x^2 & \text{if } i = n \end{cases}$ and $\beta^\epsilon P^i(x) = \begin{cases} 0 & \text{if } 2i + \epsilon > n, \epsilon = 0, 1 \\ x^p & \text{if } n = 2i, \epsilon = 0 \end{cases}$
Moreover these classes are natural with respect to homomorphisms of π -shc algebras and satisfy the Cartan formula.

Theorem B.

- 1) The algebra N^*X is naturally a π -shc algebra. Moreover, any two natural structural maps defining a π -shc structure on N^*X are π -homotopic.
- 2) If X is 1-connected then, the Jones' quasi-isomorphism $\mathfrak{C}N^*X \rightarrow C^*(X^{S^1})$ identifies the algebraic Steenrod operations, defined on HH_*N^*X by theorem A, with the topological Steenrod operations on $H^*(X^{S^1}; \mathbb{F}_p)$.

In §3 we develop the construction of a convenient model for a π -shc algebra. But, at it can be easily imagined, explicit computations with this model are not very tractable in general (cf. Example 3.9).

These two theorems are a generalization of theorem A and B, in [2]. In fact, we proved that the Jones' homomorphism induces a commutative diagram

$$\begin{array}{ccccc} N^*X & \xrightarrow{\iota} & \mathfrak{C}N^*X & \xrightarrow{\rho} & BN^*X \\ || & & J_X \downarrow & & \downarrow \overline{J}_X \\ N^*X & \xrightarrow{\iota'} & N^*X^{S^1} & \xrightarrow{\rho'} & N^*\Omega X \end{array}$$

where the horizontal arrows in the lower lines are induced from the canonical fibration $\Omega X \rightarrow X^{S^1} \rightarrow X$ and HJ_X (resp. $H\overline{J}_X$) is an isomorphism of commutative graded algebras (resp. of commutative graded Hopf algebras). Now following the lines of the proof of theorem B and of [2]-II-§3 we obtain:

Theorem C. *If X is 1-connected then the homomorphisms, $H\iota$, $H\rho$, HJ_X and $H\overline{J}_X$ respect Steenrod operations.*

The paper will be organized as follows:

§ 1 - Proof of theorem A.

§ 2 - Prof of theorem B

§ 3 - π shc model.

Appendix A - Technicalities on shc algebras.

Appendix B - Equivariant acyclic model theorem.

We would like to thank Katsuhiko Kuribayashi and Luc Menichi for helpfull conversations during the preparation of this paper.

§1 Proof of theorem A.

1.1 Preliminaries. Throughout the paper, p is a fixed prime, and we work over a field $\mathbb{F}_p$. If π is any finite group, the group ring $\mathbb{F}_p[\pi]$ is a augmented algebra. If two π -linear maps f and g are π -linear homotopic we write $f \simeq_\pi g$.

1.2. Algebraic Steenrod operations. The material involved in this section is contained in [13]. Let $\pi = \{1, \tau, \dots, \tau^{p-1}\}$ be the cyclic group of order p . We identify τ with the

p -cycle $(p, 1, 2, \dots, p-1)$ of $\mathfrak{S}_p$ thus π acts on $A^{\otimes p}$. Let $W \xrightarrow{\epsilon_W} \mathbb{F}_p$ be a projective resolution of $\mathbb{F}_p$ over $\mathbb{F}_p[\pi]$: $W = \{W_i\}_{i \geq 0}$, $\partial : W_i \rightarrow W_{i-1}$, $W_0 \cong \mathbb{F}_p[\pi]$ where each W_i is a right projective π -module and ∂ is π -linear. We choose a linear map $\eta : \mathbb{F}_p \rightarrow W$ such that $\epsilon_W \circ \eta = id_{\mathbb{F}_p}$. Necessarily, such η satisfies also $\eta \circ \epsilon_W \simeq id_W$.

Let $A = \{A_i\}_{i \in \mathbb{Z}}$ be a differential graded algebra (not necessarily associative). We denote by $m^{(p)}$ (resp. $(Hm)^{(p)}$) the iterated product $a_1 \otimes a_2 \otimes \dots \otimes a_p \mapsto a_1(a_2(\dots a_p))\dots$ (resp. the iterated product induced on HA by $m^{(p)}$). Assume that π acts trivially on A and diagonally on $W \otimes A^{\otimes p}$. If there exists a π -chain map θ such that the left hand diagram induces the right hand diagram which is commutative:

$$\begin{array}{ccccc} W \otimes A^{\otimes p} & \xrightarrow{\theta} & A & H(W \otimes A^{\otimes p}) & \xrightarrow{H\theta} & HA \\ \eta_W \otimes id \uparrow & & \nearrow m^{(p)} & \cong \uparrow & \nearrow & (Hm)^{(p)} \\ & & A^{\otimes p} & (HA)^{\otimes p} & & \end{array}$$

then for any $i \in \mathbb{Z}$ and any $x \in H^n A$ there exist well defined homology classes

$$P^i(x) \in \begin{cases} H^{n+i} A & \text{if } p = 2 \\ H^{n+2i(p-1)} A & \text{if } p > 2 \end{cases}, \quad \text{and if } p > 2, \quad \bar{P}^i(x) \in H^{n+2i(p-1)+1} A,$$

such that:

- 1) $P^i(1_{HA}) = 0, i \neq 0$,
- 2) if $p = 2$, $P^i(x) = 0$ if $i > n$ and $P^n(x) = x^2$,
- 3) if $p > 2$, $P^i(x) = 0$ if $2i > n$ and $P^i(x) = x^p$ if $n = 2i$, and $\bar{P}^i(x) = 0$ if $2i \geq n$,
- 4) If A is the mod p reduction of a free graded module over $\mathbb{Z}$ then $\beta \circ P^{i-1} = iP^i$ if

$p = 2$ and if $p > 2$ $\bar{P}^i = \beta \circ P^i$.

Moreover these classes do not depend on the choice of W neither on η , and are natural with respect to homomorphisms of algebras commuting with the structural maps θ . These *algebraic Steenrod operations* do not in general satisfy: $P^i(x) = 0$ if $i < 0$, $P^0(x) = x$, the Cartan formula and the Adem relations.

1.3 Cartan formula. Assume $A^i = 0, i < 0$. The structural map $\theta : W \otimes A^{\otimes p} \rightarrow A$ induces $\tilde{\theta} : A^{\otimes p} \rightarrow \text{Hom}(W, A)$, $\tilde{\theta}(u)(w) = (-1)^{|u||w|} \theta(w \otimes u)$, $u \in A^{\otimes p}, w \in W$. Let us precise that

if $f \in \text{Hom}^k(W, A) = \prod_{i \geq 0} \text{Hom}(W_i, A^{k-i}) = \bigoplus_{i=0}^k \text{Hom}(W_i, A^{k-i})$ then: $Df = d \circ f - (-1)^k f \circ \partial$,

$(\sigma f)(w) = f(w\sigma)$, $\sigma \in \pi, w \in W$. A direct verification shows that $\text{Hom}^k(W, A)$ is a π -complex and that $\tilde{\theta}$ is a π -chain map. If $ev_0 : \text{Hom}(W, A) \rightarrow A$ denotes the evaluation map on the generator $e_0 = \eta(1)$ of W_0 , the left hand diagram induces the right hand diagram which is commutative.

$$\begin{array}{ccccc} A^{\otimes p} & \xrightarrow{\tilde{\theta}} & \text{Hom}(W, A) & (HA)^{\otimes p} & \xrightarrow{H\tilde{\theta}} & H(\text{Hom}(W, A)) \\ m^{(p)} \searrow & & \downarrow ev_0 & (Hm)^{(p)} \searrow & & \downarrow H(ev_0) \\ & & A & & & HA \end{array}$$

Let $\psi : W \rightarrow W \otimes W$ be a diagonal approximation and let m denotes the product in the algebra A . The formula $f \cup g = m \circ (f \otimes g) \circ \psi$ defines cup product $\text{Hom}^k(W, A) \otimes \text{Hom}^l(W, A) \rightarrow \text{Hom}^{k+l}(W, A)$, $f \otimes g \mapsto f \cup g$ and $\text{Hom}(W, A)$ is (non associative) differential graded algebra.

1.4 Proposition. Let $(A, \tilde{\theta})$ be as above. If we assume that $H^* \tilde{\theta}$ respects products then the algebraic Steenrod operations defined by $\tilde{\theta}$ satisfy the Cartan formula: $P^i(xy) = \sum_{j+k=i} P^j(x) P^k(y)$, $x, y \in H^* A$.

Proof. We consider the *standard small free resolution* of a finite cyclic group [5]: the (right) π -free acyclic chain complex defined by: $W = \{W_i\}_{i \geq 0}$, $W_i = e_i \mathbb{F}_p[\pi]$, $\partial : W_i \rightarrow W_{i-1}$ and if τ is a fixed generator of π then $\partial e_{2i+1} = (1 + \tau)e_{2i}$, $\partial e_{2i} = (1 + \tau + \dots + \tau^{p-1})e_{2i-1}$. The augmentation $\epsilon_W : W \rightarrow \mathbb{F}_p$ is defined by $\epsilon_W(e_i) = 0, i > 0$ and $\epsilon_W(e_0) = 1$. Let $\Delta A \subset A^{\otimes p}$ be the subspace generated by the image of the diagonal map $\Delta : A \rightarrow A^{\otimes p}$ $a \mapsto a^{\otimes p}$. Thus π acts trivially on the sub-algebra ΔA , and there exists a π -module $M \subset A^{\otimes p}$ such that, [13]-Lemma 1.3, $H(W \otimes_\pi A^{\otimes p}) = (\bigoplus_{i=0}^\infty \mathbb{F}_p e_i \otimes \Delta A) \oplus (\mathbb{F}_p e_0 \otimes M)$. The algebraic Steenrod operations are defined by ([13]-Proposition 2.3): for $x \in H^n A$ if $p = 2$ by, $P^i(x) = \theta^*(e_{n-i} \otimes x^{\otimes p}) := Sq^i(x)$ and if p is odd by, $P^i(x) = (-1)^i \nu(n) \theta^*(e_{(n-2i)(p-1)} \otimes x^{\otimes p})$, $\overline{P}^i(x) = (-1)^i \nu(n) \theta^*(e_{(n-2i)(p-1)-1} \otimes x^{\otimes p})$. Here $\nu(n) = (-1)^j \left(\left(\frac{p-1}{2}\right)!\right)^\epsilon$ if $n = 2j + \epsilon$, $\epsilon = 0, 1$ and θ^* denotes the map induced by θ on $H(W \otimes_\pi A^{\otimes p})$. A diagonal approximation is explicitly given by, [4] -§31-Chap. XII: $\psi e_i(\tau^s) = \psi(e_i) \tau^s$, $\psi(e_{2l+1}) = \sum_{j+k=l} e_{2j} \otimes e_{2k+1} + \sum_{j+k=l} e_{2j+1} \otimes e_{2k} \tau^{-1}$ and $\psi(e_{2l}) = \sum_{j+k=l} e_{2j} \otimes e_{2k} + \sum_{j+k=l-1} \sum_{0 \leq r < s < p} e_{2j+1} \tau^{-r} \otimes e_{2k+1} \tau^{-s}$. For simplicity, let us assume that $p = 2$. First observe that the restriction of $\tilde{\theta}$ on $\Delta A \subset A^{\otimes 2}$ factorizes through a map $\tilde{\theta}_\pi : \Delta A \rightarrow \text{Hom}_\pi(W, A)$ where $\text{Hom}_\pi(W, A)$ denotes the sub-complex of π -linear maps. Secondly, note that $H^*(\text{Hom}_\pi(W, A)) \cong H^*(\text{Hom}_\pi(W, H^* A))$ and that the map $H^* \tilde{\theta}$ factorizes through a map $(H^* \tilde{\theta})_\pi : K \cap \ker d \rightarrow \text{Hom}_\pi(W, H^* A)$. Let us denote by $j : \text{Hom}_\pi(W, A) \rightarrow \text{Hom}(W, A)$ the natural inclusion of complexes. By definition, if $x \in H^n A$ and a denotes any cocycle such that $x = cl(a)$: $Sq^i x = H^* j \circ ((H^* \tilde{\theta})_\pi(a^{\otimes 2}))(e_{n-i}) = cl(\tilde{\theta}(a^{\otimes 2})(e_{n-i}))$. If $y = cl(b) \in H^m A$ then:

$$\begin{aligned} Sq^i(xy) &= cl(\tilde{\theta}(ab \otimes ab)(e_{n+m-i})) = cl(\tilde{\theta}((a \otimes a)(b \otimes b))(e_{n+m-i})) \\ &= cl((\tilde{\theta}(a \otimes a) \cup \tilde{\theta}(b \otimes b))(e_{n+m-i})) = cl\left(\sum_{j=0}^{n+m-i} \tilde{\theta}(a \otimes a)(e_j) \tilde{\theta}(b \otimes b)(e_{m+n-i-j} \tau^j)\right) \\ &= \sum_{j=0}^{n+m-i} cl(\tilde{\theta}(a \otimes a)(e_j)) cl(\tilde{\theta}(b \otimes b)(e_{m+n-i-j})) = \sum_{j=0}^{n+m-i} Sq^{n-j} x Sq^{-n+i+k} y \\ &= \sum_{j+k=i} Sq^j x Sq^k y. \end{aligned}$$

1.5. The differential graded algebra $\mathfrak{C}_*(A)$. The bar and the cobar construction can be regarded as a pair of adjoint functors $B : \mathbf{DA} \leftrightarrow \mathcal{N}\text{-}\mathbf{DC} : \Omega$, where $\mathbf{DA}$ is the category of supplemented differential graded associative algebras and $\mathcal{N}\text{-}\mathbf{DC}$ is the category of locally conilpotent supplemented graded coassociative coalgebras. (See for instance, [6]-2.14 for more details.) This leads to a natural homomorphism $\alpha_A : \Omega BA \rightarrow A$ of differential graded algebras. The elements of BA (resp. ΩC) are denoted $[a_1|a_2|\dots|a_k] \in B_k A$, and $[\] := 1 \in B_0 A \cong \mathbb{F}_p$ (resp. $\langle c_1|c_2|\dots|c_k \rangle \in \Omega_k A$, and $\langle \ \rangle := 1 \in \Omega_0 A \cong \mathbb{F}_p$). We remind the reader that the linear map $i_A : A = \mathbb{F}_p \oplus \overline{A} \rightarrow \Omega BA$, $i_A(1) = 1$ and $i_A(a) = s^{-1}sa = \langle a \rangle, a \in \overline{A}$, commutes with the differentials and satisfies: $\alpha_A \circ i_A = id$, $i_A \circ \alpha_A \simeq id_{\Omega BA}$.

Recall also that the *Hochschild complex* of A , is the graded vector space $\mathfrak{C}_* A = \{\mathfrak{C}_k A\}_{k \geq 0}$, $\mathfrak{C}_k A = A \otimes B_k A$, where a generator of $\mathfrak{C}_k A$ is of the form $a_0[a_1|a_2|\dots|a_k]$ if $k > 0$ and $a[\]$ if $k = 0$. We set $\epsilon_i = |a_0| + |sa_1| + |sa_2| + \dots + |sa_{i-1}|$, $i \geq 1$ and define the differential $d = d^1 + d^2$ by:

$$\begin{aligned} d^1 a_0[a_1|a_2|\dots|a_k] &= da_0[a_1|a_2|\dots|a_k] - \sum_{i=1}^k (-1)^{\epsilon_i} a_0[a_1|\dots|da_i|\dots|a_k] \\ d^2 a_0[a_1|a_2|\dots|a_k] &= (-1)^{|a_0|} a_0 a_1[a_2|\dots|a_k] + \sum_{i=2}^k (-1)^{\epsilon_i} a_0[a_1|\dots|a_{i-1}a_i|\dots|a_k] \\ &\quad - (-1)^{|sa_k| \epsilon_k} a_k a_0[a_1|\dots|a_{k-1}] \end{aligned}$$

Henceforth we shall discard the lower star in $\mathfrak{C}_* A$ and write $\mathfrak{C}A$. The associativity of the algebra A is necessary to prove that $d \circ d = 0$ in BA as well as in $\mathfrak{C}A$ and if A is commutative then BA (resp. $\mathfrak{C}A$) is a differential graded Hopf algebra (resp. a differential graded algebra.) Since we are interested to the case $A = N^*(X)$, and since this differential

graded algebra is not commutative but only *strongly homotopy commutative* (a *shc* algebra for short), [15], it seems natural for our purpose, to consider *shc* algebra in order to define a product on $\mathfrak{C}A$.

A *shc algebra* is a pair (A, μ) such that $\mu : \Omega B(A^{\otimes 2}) \rightarrow \Omega BA$ is a homomorphism of differential graded algebras where $\alpha \circ \mu \circ i_{A \otimes A} = m_A$ (m_A denotes the product in A). Moreover, μ satisfies the unity axiom, the associativity axiom and the commutativity axiom as described in [15]-4.1. Recall that two homomorphisms of differential graded algebras $f, g : A \rightarrow A'$ are *homotopic* (we write $f \simeq_{\mathbf{DA}} g$, even if A and A' are not associative) if there exists a (f, g) -derivation h such that $f - g = dh + hd$. Let (A, d, μ) and (A', d', μ') be two *shc* algebras. A *strict shc homomorphism* $f : (A, d, \mu) \rightarrow (A', d', \mu')$ is a morphism $f \in \mathbf{DA}(A, A')$ such that $\Omega Bf \circ \mu \simeq_{\mathbf{DA}} \mu' \circ \Omega B(f \otimes f)$.

The chain map $\mathfrak{C}(\alpha_A)$ is surjective and $H\mathfrak{C}(\alpha_A) = HH_*(\alpha_A)$ is an isomorphism. We denote by $s_A : \mathfrak{C}A \rightarrow \mathfrak{C}\Omega BA$ any chain map such that $\mathfrak{C}(\alpha_A) \circ s_A = id$ and $s_A \circ \mathfrak{C}(\alpha_A) \simeq id$. When (A, μ) is a *shc* algebra a product $m_{\mathfrak{C}A} : \mathfrak{C}A \otimes \mathfrak{C}A \rightarrow \mathfrak{C}A$ is defined by the composite

$$\mathfrak{C}A \otimes \mathfrak{C}A \xrightarrow{sh} \mathfrak{C}(A \otimes A) \xrightarrow{s_{A \otimes A}} \mathfrak{C}\Omega B(A \otimes A) \xrightarrow{\mathfrak{C}\mu_A} \mathfrak{C}\Omega BA \xrightarrow{\mathfrak{C}\alpha_A} \mathfrak{C}A$$

where *sh* denotes the shuffle product, [2]. More precisely, we proved: *If A is a shc algebra then*

- 1) *BA is a differential graded Hopf algebra such that HBA is a commutative graded Hopf algebra,*
- 2) *$\mathfrak{C}A$ is a (non associative) differential graded algebra such that $HH_*(A) := H\mathfrak{C}A$ is a commutative graded algebra,*
- 3) *If A' is an other shc algebra and if f is a homomorphism of shc algebras then there is a commutative diagram*

$$\begin{array}{ccccc} A & \xrightarrow{\iota} & \mathfrak{C}A & \xrightarrow{\rho} & BA \\ f \downarrow & & \mathfrak{C}f \downarrow & & \downarrow Bf \\ A' & \xrightarrow{\iota'} & \mathfrak{C}A' & \xrightarrow{\rho'} & BA' \end{array}$$

where the maps $\iota, \iota', \rho, \rho'$ and $\mathfrak{C}f$ are homomorphisms of differential graded algebras, Bf is a homomorphism of differential graded Hopf algebras.

If A is a *shc* algebra and if $p = 2$, then both Sq^n and Sq^{n-1} are determined on $H^n A$, ([2]-6.6 or [15]-4.8), while Sq^{n-2} is not determined on $(HH_* A)^n$, (cf. Example 3.8). Therefore, we need to enrich the notion of *shc* algebra.

1.6 π -shc algebra. Let (A, d, μ) be a *shc* algebra. For any $n \geq 2$, by Lemma A.3, there exists a homomorphism of differential graded algebras, called the *shc iterated structural map* $\mu^{(n)} : \Omega B(A^{\otimes n}) \rightarrow \Omega BA$ such that: $\mu^{(2)} = \mu$ and $\alpha_A \circ \mu^{(n)} \circ i_{A^{\otimes n}} \simeq m_A^{(n)}$. Let W, ψ as in 1.3.

A *shc* algebra (A, μ) is, a π -*shc algebra* if there exists a map $\tilde{\kappa}_A : \Omega B(A^{\otimes p}) \rightarrow \text{Hom}(W, A)$ which is both a π -linear map and a homomorphism of differential graded algebras such that the next diagram commutes, up to a derivation homotopy.

$$\begin{array}{ccc} \Omega B(A^{\otimes p}) & \xrightarrow{\tilde{\kappa}_A} & \text{Hom}(W, A) \\ \alpha_A \circ \mu^{(p)} \searrow & & \downarrow ev_0 \\ & & A \end{array} .$$

Here $\sigma \in \mathfrak{S}_p$ acts on $B(A^{\otimes p})$ by the rule $\sigma[x_1|x_2|\dots|x_k] = [\sigma x_1|\sigma x_2|\dots|\sigma x_k]$, $x_i \in A^{\otimes p}$ and on $\Omega B(A^{\otimes p})$ by the rule $\sigma \in \pi$, $\sigma < y_1|y_2|\dots|y_l > = < \sigma y_1|\sigma y_2|\dots|\sigma y_l >$, $y_j \in B(A^{\otimes p})$.

A *strict π -shc homomorphism* $f : (A, \mu_A, \tilde{\kappa}_A) \rightarrow (A', \mu_{A'}, \tilde{\kappa}_{A'})$ is a strict *shc* homomor-

phism satisfying $\kappa_{A'} \circ \Omega B(f^{\otimes p}) \simeq_{\pi\text{-DA}} \text{Hom}(W, f) \circ \tilde{\kappa}_A$.

If $(A, \mu, \tilde{\kappa}_A)$ is a π -shc algebra then the next diagram commutes, up to a π -linear homotopy:

$$\begin{array}{ccc} \Omega B(A^{\otimes p}) & \xrightarrow{\tilde{\kappa}_A} & \text{Hom}(W, A) \\ \alpha_{A^{\otimes p}} \downarrow & & \downarrow ev_0 \\ A^{\otimes p} & \xrightarrow{m^{(p)}} & A \end{array},$$

and the Steenrod operations, defined for $x = cl(a) \in H^n A$ by : $Sq^i(x) = cl(\tilde{\kappa}(i_{A^{\otimes 2}}(a^{\otimes 2}))(e_{n-i}))$ if $p = 2$ and by $P^i(x) = (-1)^i \nu(n) cl(\tilde{\kappa}(i_{A^{\otimes p}}(a^{\otimes p}))(e_{(n-2i)(p-1)}))$ if $p > 2$, satisfy the Cartan formula (see Proof of 1.4 for notations).

In order to construct Steenrod operations on $\mathfrak{C}(A)$ we need to compare $\text{Hom}(W, \mathfrak{C}A)$ and $\mathfrak{C}\text{Hom}(W, A)$. But this last expression is not defined unless $\text{Hom}(W, A)$ is associative. If $p \neq 2$, the diagonal approximation considered in the proof of Proposition 1.4 is not coassociative. Thus, for p odd, we will consider the *standard resolution* of a finite cyclic group π , [14], [1]. The Alexander-Whitney diagonal approximation ψ_W , admits the counity ϵ and is strictly coassociative.

1.7. Proposition. *If A is a differential graded algebra and if (W, ψ_W) is a coassociative coalgebra then there exists a natural chain map ϕ_A such that the following diagram commutes*

$$\begin{array}{ccc} \mathfrak{C}\text{Hom}(W, A) & \xrightarrow{\phi_A} & \text{Hom}(W, \mathfrak{C}A) \\ \mathfrak{C}ev_0 \searrow & & \swarrow ev_0 \\ & \mathfrak{C}A & \end{array}$$

Moreover, ϕ is π -linear (π acts trivially on $B = A, \mathfrak{C}A$, diagonally on $\text{Hom}(W, B)$).

Proof. We define ϕ_A by: $\phi_A(f_0[]) = f_0$ and if $k > 0$ by $\phi_A(f_0[f_1|f_2|\dots|f_k]) = (id \otimes s^{\otimes k}) \circ (f_0 \otimes f_1 \otimes f_2 \otimes \dots \otimes f_k) \circ \psi_W^{(k)}$ where $\psi_W^{(k)}$ denotes the iterated diagonal: $\psi_W^{(1)} = \psi_W$, $\psi_W^{(k+1)} = (id \otimes \psi_W) \circ \psi_W^{(k)}$ and $s : A \rightarrow sA$ denotes the suspension of degrees. Obviously, ϕ_A is natural in A . Let us check in detail that ϕ_A commutes with the differentials. (σ_k denotes the cycle $(1, \dots, k) \rightarrow (k, 1, \dots, k-1)$)

$$\begin{aligned} \phi_A(d^1(f_0[f_1|f_2|\dots|f_k])) &= \phi_A(df_0[f_1|f_2|\dots|f_k] - \sum_{i=1}^k (-1)^{\epsilon_i} f_0[f_1|\dots|df_i|\dots|f_k]) \\ &= (id \otimes s^{\otimes k}) \left(df_0 \otimes f_1 \otimes f_2 \otimes \dots \otimes f_k - \sum_{i=1}^k (-1)^{\epsilon_i} f_0 \otimes f_1 \otimes \dots \otimes df_i \otimes \dots \otimes f_k \right) \circ \psi_W^{(k)} \\ &= (id \otimes s^{\otimes k}) \left((d_A \circ f_0 - (-1)^{|f_0|} f_0 \circ \partial) \otimes f_1 \otimes f_2 \otimes \dots \otimes f_k \right. \\ &\quad \left. - \sum_{i=1}^k (-1)^{\epsilon_i} f_0 \otimes f_1 \otimes \dots \otimes (d_A \circ f_i - (-1)^{|f_i|} f_i \circ \partial) \otimes \dots \otimes f_k \right) \circ \psi_W^{(k)} \\ &= (id \otimes s^{\otimes k}) \left((d_A \otimes id - id \otimes d_{A^{\otimes k}}) \circ (f_0 \otimes f_1 \otimes f_2 \otimes \dots \otimes f_k) \circ \psi_W^{(k)} \right. \\ &\quad \left. - (-1)^{|f_0|+|f_1|+\dots+|f_k|+k} (id \otimes s^{\otimes k}) \circ (f_0 \otimes f_1 \otimes \dots \otimes f_i \otimes \dots \otimes f_k) \circ \partial_{W^{\otimes k}} \right) \circ \psi_W^{(k)} \\ &= d_{\mathfrak{C}A}^1 \circ \phi_A((f_0[f_1|f_2|\dots|f_k]) - (-1)^{\epsilon_k} \phi_A((f_0[f_1|f_2|\dots|f_k]) \circ \partial) \\ \phi_A(d^2(f_0[f_1|f_2|\dots|f_k])) &= \phi_A \left((-1)^{|f_0|} (f_0 \cup f_1)[f_2|\dots|f_k] + \sum_{i=2}^k (-1)^{\epsilon_i} f_0[f_1|\dots|f_i \cup f_i|\dots|f_k] \right. \\ &\quad \left. + (-1)^{|s f_k| \epsilon_k} (f_k \cup f_0)[f_1|\dots|f_{k-1}] \right) \\ &= (id \otimes s^{\otimes k-1}) \circ \left((-1)^{|f_0|} (m_A \circ (f_0 \otimes f_1) \circ \psi_W) \circ (f_2 \otimes \dots \otimes f_k) \right) \end{aligned}$$

$$\begin{aligned}
& + \sum_{i=2}^k (-1)^{\epsilon_i} f_0[f_1|\dots|m_A \circ (f_i \otimes f_i) \circ \psi_W|\dots|f_k] + \\
& \quad (-1)^{|sf_k|\epsilon_k} (m_A \circ (f_k \otimes f_0) \circ \psi_W)[f_1|\dots|f_{k-1}] \circ \psi_W^{(k-1)} \\
& = (id \otimes s^{\otimes k-1}) \circ \left((m_A \otimes id + \sum_{i=2}^k (-1)^i id \otimes m_A \otimes id + (-1)^{k+1} (m_A \otimes id) \circ \sigma_k) \right. \\
& \quad \left. \circ (1 \otimes s^{\otimes k})^{-1} \circ \phi_A((f_0[f_1|f_2|\dots|f_k])) \right) \\
& = d_{\mathfrak{C}A}^2 \phi_A(f_0[f_1|f_2|\dots|f_k]).
\end{aligned}$$

Observe that we need also the coassociativity property of the coproduct ψ_W in the proof of the second equality. It is straightforward to check that the remaining claims in 1.7 hold.

1.8 End of the proof of Theorem A. One also define, for $n \geq 3$, $(k_1, k_2, \dots, k_n)$ -shuffles, the n -iterated shuffle map $sh^{(n)} : (\mathfrak{C}A)^{\otimes n} \rightarrow \mathfrak{C}(A^{\otimes n})$ and the iterated product $m_{\mathfrak{C}A}^{(p)} : \mathfrak{C}A^{\otimes p} \rightarrow \mathfrak{C}A$:

$$\mathfrak{C}A^{\otimes p} \xrightarrow{sh^{(p)}} \mathfrak{C}(A^{\otimes p}) \xrightarrow{s_{A^{\otimes p}}} \mathfrak{C}\Omega B(A^{\otimes p}) \xrightarrow{\mathfrak{C}\mu_A^{(p)}} \mathfrak{C}\Omega BA \xrightarrow{\mathfrak{C}\alpha_A} \mathfrak{C}A, \quad (\text{compare with 1.5}).$$

Consider ϕ_A as defined above, then applying the functor $\mathfrak{C}$ to the diagram that appear in the definition of a π -shc algebra (1.6), we obtain the following diagram:

$$\begin{array}{ccc}
\mathfrak{C}\Omega B(A^{\otimes p}) & \xrightarrow{\phi_A \circ \mathfrak{C}\tilde{\kappa}_A} & \text{Hom}(W, \mathfrak{C}A) \\
\mathfrak{C}(\alpha_{A^{\otimes p}}) \downarrow & & \downarrow ev_0 \\
\mathfrak{C}(A^{\otimes p}) & & \\
sh^{(p)} \uparrow & & \\
(\mathfrak{C}A)^{\otimes p} & \xrightarrow{m_{\mathfrak{C}A}^{(p)}} & \mathfrak{C}A.
\end{array}$$

Since $ev_0 \circ \tilde{\kappa}_A \simeq_{\mathbf{DA}} \alpha_A \circ \mu_A^{(p)}$, Lemma A.6 implies that $ev_0 \circ \phi_A \circ \mathfrak{C}\tilde{\kappa}_A \simeq \mathfrak{C}\mu^{(p)}$. Now observe that in the sequence $\mathfrak{C}\Omega B(A^{\otimes p}) \xrightarrow{\mathfrak{C}(\alpha_{A^{\otimes p}})} \mathfrak{C}(A^{\otimes p}) \xleftarrow{(sh^{(p)})} (\mathfrak{C}A)^{\otimes p}$ of chain maps:

a) $\mathfrak{C}(\alpha_{A^{\otimes p}})$ is π -linear. Indeed $\sigma \in \mathfrak{S}_p$ acts on $\mathfrak{C}(A^{\otimes p})$ (resp. on $\mathfrak{C}\Omega B(A^{\otimes p})$) by the rule $\sigma(x_0[x_1|x_2|\dots|x_k]) = \sigma x_0[\sigma x_1|\sigma x_2|\dots|\sigma x_k]$, $x_i \in A^{\otimes p}$ (resp. $x_i \in \Omega BA^{\otimes p}$). Then, $\mathfrak{C}\Omega B(A^{\otimes p})$ and $\mathfrak{C}(A^{\otimes p})$ are $\mathfrak{S}_p$ -complexes, $\alpha_{A^{\otimes p}}$ and thus $\mathfrak{C}\alpha_{A^{\otimes p}}$ are $\mathfrak{S}_p$ -linear.

b) $sh^{(p)}$ is also $\mathfrak{S}_p$ -linear. Indeed, one simply writes: $sh^{(p)}(a_0^1[a_1^1|a_2^1|\dots|a_{n_1}^1] \otimes a_0^2[a_1^2|\dots|a_{n_2}^2] \otimes \dots \otimes a_0^p[a_1^p|\dots|a_{n_p}^p]) = (a_0^1 \otimes a_0^2 \otimes \dots \otimes a_0^p) \sum_{\sigma} \sigma[a_1^1 \otimes 1 \otimes 1 \otimes \dots \otimes 1 | \dots | a_{n_1}^1 \otimes 1 \otimes 1 \otimes \dots \otimes 1 | 1 \otimes a_1^2 \otimes 1 \otimes \dots \otimes 1 | \dots | 1 \otimes 1 \otimes 1 \otimes \dots \otimes 1 \otimes a_i^p | \dots | 1 \otimes 1 \otimes 1 \otimes \dots \otimes 1 \otimes a_{n_p}^p]$ where σ ranges over all $(n_1, n_2, \dots, n_p)$ -shuffles and observe that the set of all $(n_1, n_2, \dots, n_p)$ -shuffles coincides with the set of all $(\lambda(n_1), \lambda(n_2), \dots, \lambda(n_p))$ -shuffles when $\lambda \in \mathfrak{S}_p$.

c) $\mathfrak{C}(\alpha_{A^{\otimes p}})$ is a homomorphism of differential graded algebras. This follows directly from F4, F6 and F8.

From, properties a), b), c) and following the lines of the proof on [13]-Proposition 2.3, (sketched in the proof of Proposition 1.4), using the section $s_{A^{\otimes p}}$, this diagram defines natural Steenrod operations on $HH_*(A)$. In order to establish the Cartan formula it is enough, by Proposition 1.4, to prove that $H^*\tilde{\kappa}_A$ respects products. This is Lemma A.5. Naturality follows directly from Lemma A.6.

§2 - Proof of theorem B.

2.1 Twisting cochains. In [15]-4.7, H.J. Munkholm has established the existence of a natural transformation $\mu_X : \Omega B(N^*X \otimes N^*X) \rightarrow \Omega B N^*X$ such that $\alpha_{N^*X} \circ \mu_X \circ i_{N^*X \otimes N^*X}$ is the usual cup product $N^*X \otimes N^*X \rightarrow N^*X$. Moreover, (N^*X, μ_X) is a shc-algebra. Let us denote by $\mu_X^{(p)} : \Omega B((N^*X)^{\otimes p}) \rightarrow \Omega B N^*X$ the iterated shc structural map and by

t_X the twisting cochain associated to $\alpha_{N^*X} \circ \mu_X^{(p)}$: $t_X \in \text{Hom}^1(B((N^*X)^{\otimes p}), N^*X)$, $t \cup t = Dt$, $t(x)(e_0) = \mu(< x >)$, $x \in \overline{BN^*X}$, $t([]) = 0$ where D denotes the differential in $\text{Hom}(B((N^*X)^{\otimes p}), N^*X)$ (See [15]-1.8 for more details on twisting cochains). We want to construct a twisting cochain $t'_X \in \text{Hom}^1(B((N^*X)^{\otimes p}), \text{Hom}(W, N^*X))$, $t' \cup t' = D't'$ such that: $ev_0 \circ t'_X = t_X$, t'_X is natural in X and t'_X is π -linear. (Here D' denotes the differential in $\text{Hom}(B((N^*X)^{\otimes p}), \text{Hom}(W, N^*X))$.) Such a twisting cochain t'_X will determine a homomorphism of differential graded algebras $\tilde{\kappa}_X : \Omega B(N^*X)^{\otimes p} \rightarrow \text{Hom}(W, N^*X)$ such that $(N^*X, \mu_X, \tilde{\kappa}_X)$ is a π -shc algebra. Moreover, by Corollary B.4, $\tilde{\kappa}_X$ is well determined up to a π -linear homotopy, and so the first part of theorem B will be proved.

2.2 Proof of part 1. We construct t' by induction on the degree in $W = \{W_i\}_{i \geq 0}$ and on the bar degree: $\overline{B((N^*X)^{\otimes p})} = \{B_k((N^*X)^{\otimes p})\}_{k \geq 1}$. We denote by Δ the coproduct in $B((N^*X)^{\otimes p})$. For simplicity, if $x \in B_k(N^*X)^{\otimes p}$ we write $\Delta x = 1 \otimes x + x \otimes 1 + \overline{\Delta}x = x' \otimes x''$ where $\overline{\Delta}x \in \oplus_{l'+l''=k} B_{l'}((N^*X)^{\otimes p}) \otimes B_{l''}((N^*X)^{\otimes p})$ with l' (resp l'') such that $0 < l' < k$ (resp. $0 < l'' < k$).

First, if e_0 denotes a generator of W_0 , we define t' on $W_0 \otimes \overline{B((N^*X)^{\otimes p})}$ by setting:

$$(\star_0) \quad t'(x)(e_0\sigma) = t(\sigma^{-1}x), \quad \sigma \in \pi, \quad x \in B_k N^*X.$$

t' is a π -linear map, that is: if $\sigma \in \pi$ and $x \in \overline{B(N^*X)^{\otimes p}}$ then $t'(\sigma x) = \sigma(t'(x))$. It suffices to prove this last formula for the elements $e_0\tau^j$ in W_0 :

$$\begin{aligned} (\sigma t'(x))(e_0\tau^j) &= t'(x)(e_0\tau^j\sigma^{-1}) = t'(x)(e_0(\sigma(\tau^j)^{-1})^{-1}) \\ &= t'(\sigma(\tau^j)^{-1}x)(e_0) = t'((\tau^j)^{-1}\sigma x)(e_0) = t'(\sigma x)(e_0\tau^j). \end{aligned}$$

Precise that π acts diagonally on $B(N^*X)^{\otimes p} \otimes B(N^*X)^{\otimes p}$ and on $W \otimes W$ and that Δ (resp. ψ) are π -linear. We check in detail that: $D't'(x)(w) = (t' \cup t')(x)(w)$, $w \in W_0$, $x \in \overline{B(N^*X)^{\otimes p}}$. Once again, it suffices to prove this formula for the elements $e_0\sigma \in W_0$:

$$\begin{aligned} D't'(x)(e_0\sigma) &= Dt'(x)(e_0\sigma) + t'(dx)(e_0\sigma) = d(t'(x)(e_0\sigma)) + t'(x)(\partial e_0\sigma) - t'(dx)(e_0\sigma) \\ &= d(t(\sigma^{-1}x)) + t(\sigma^{-1}dx) = (Dt)(\sigma^{-1}x) = (t \cup t)(\sigma^{-1}x) \\ &= (m_{N^*X} \circ (t \otimes t) \circ \Delta)(\sigma^{-1}x) = m_{N^*X} \circ (t \otimes t)(\sigma^{-1}\Delta x) \\ &= m_{N^*X}(t'(\sigma^{-1}x')(e_0) \otimes t'(\sigma^{-1}x'')(e_0)) = m_{N^*X}(t'(x')(e_0\sigma) \otimes t'(x'')(e_0\sigma)) \\ &= (m_{N^*X} \circ (t' \otimes t') \circ \Delta)(x)(\psi(e_0\sigma)) = (t' \cup t')(x)(e_0\sigma). \end{aligned}$$

Secondly, for an arbitrary generator $e_i \in W_i$, $i \geq 1$ and for $[x] \in B_1((N^*X)^{\otimes p})$, we define

$$(\star_1) \quad t'([x])(e_i\sigma) = \begin{cases} 0 & \text{if } x \in V_k \\ \lambda(\sigma^{-1}x)(e_i) & \text{if not} \end{cases}, \quad \text{where } V_k = \mathbb{F}_p^{\otimes p-k-1} \otimes N^*X \otimes \mathbb{F}_p^{\otimes p-k} \text{ and } \lambda$$

is a natural π -linear transformation of degree 0 such that the following diagram commutes up to homotopy

$$\begin{array}{ccc} (N^*X)^{\otimes p} & \xrightarrow{\lambda_X} & \text{Hom}(W, N^*X) \\ & m_{N^*X}^{(p)} \searrow & \downarrow ev_0 \\ & & N^*X. \end{array}$$

Such a transformation exists for Corollary B.4 applies. Observe that if $w \in W_i$ and $x \in ((N^*X)^{\otimes p})$ then $(d_{N^*X}(\lambda_X(x)) - \lambda_X(dx))(w) = (-1)^{|x|}\lambda(x)(\partial w)$. The formula $(\star_1)$ defines a π -linear map $t' : B_1((N^*X)^{\otimes p}) \rightarrow \text{Hom}(W, N^*X)$ and since $\overline{\Delta}[x] = 0$ it follows that $Dt'([x]) = 0 = (t' \cup t')[x]$, $[x] \in B_1((N^*X)^{\otimes p})$.

Thirdly, we assume that, for some $k \geq 1$, there is a π -linear map $t' : \overline{B_{<k}((N^*X)^{\otimes p})} \rightarrow \text{Hom}(W, N^*X)$, which extends to a π -linear map $t' : B_k((N^*X)^{\otimes p}) \rightarrow \text{Hom}(W_{<i}, N^*X)$, for some $i \geq 1$ and satisfying $D't'(y) = (t' \cup t')(y)$, $y \in \overline{B_{<k}((N^*X)^{\otimes p})}$. Let d_1 be the internal part of the differential in the bar construction $d = d_1 + d_2$. The contravariant

functor $X \mapsto FX = (\mathbb{F}_p \oplus s^i B_k((N^*X)^{\otimes p}), d)$ where $d(s^i y) = s^i dy$, $y \in B_k((N^*X)^{\otimes p})$ is acyclic on the models $\{\Delta^n\}_{n \geq 0}$. We define the natural transformation: $\mathcal{T}_{i,X} : FX \rightarrow N^*X$, $y \mapsto \mathcal{T}_{i,X}(y) = (t' \cup t')(y)(e_i) - (-1)^{|y|} t'(y)(\partial e_i) - t'(d^2 y)(e_i)$. (The inductive hypothesis ensures us that $\mathcal{T}_{i,X}(y)$ is well defined.) Let us check in detail that $\tilde{D}\mathcal{T}_{i,X} = 0$ when $\tilde{D}$ denotes the differential in $\text{Hom}(FX, N^*X)$. Observe that, for any generator e_i of W_i and if $y \in FX$, the inductive hypothesis implies : $(D't')(y)(\partial e_i) = (t' \cup t')(\partial e_i)$ and $(D't')(d_2 y)(e_i) = (t' \cup t')(d_2 y)(e_i)$. Thus we obtain:

$$\begin{aligned} (\tilde{D}\mathcal{T}_{i,X})(y) &= d_{N^*X}((t' \cup t')(y)(e_i) - (-1)^{|y|} (D't')(y)(\partial e_i) \\ &\quad + (-1)^{|t'(y)|} t'(y)(\partial^2 e_i) + (-1)^{|t'(y)|} t'((d_1 + d_2)(y))(\partial e_i) \\ &\quad - ((D't')(d_2 y)(e_i) + (-1)^{|t'(D_2 y)|} t'(d_2 y)(\partial e_i) + (-1)^{|t'|} t'((d_1 + d_2)(d_2 y)(e_i)) \\ &\quad - ((t' \cup t')(d_1 y)(e_i) - (-1)^{|y|+1} t'(d_1 y)(\partial e_i) - t'(D_2 D_1 y)(e_i))) \\ &= D'(t' \cup t')(y)(e_i) + (-1)^{|y|} (t'(d_1 + d_2)(y)(\partial e_i) - t'(d_1 y)(\partial e_i) - t'(d_2 y)(\partial e_i)) \\ &\quad + (t'(d_1 \circ d_2 y)(e_i) + t'(d_2 \circ d_1 y)(e_i)). \end{aligned}$$

Now $(\tilde{D}\mathcal{T}_{i,X})(y) = 0$, since each line in the last equality is 0.

Theorem B.4, establishes the existence of a natutal transformation $\mathcal{S}_{i,X} : FX \rightarrow N^*X$ such that $D'\mathcal{S}_{i,X} = \mathcal{T}_{i,X}$ that is: if $z \in FX$ then $\mathcal{T}_{i,X}(z) = d\mathcal{S}_{i,X}(z) - (-1)^{|\mathcal{S}_{i,X}|} d^1 z$. We set for $y \in B_k((N^*X)^{\otimes p})$, $t'(y)(e_i \sigma) = \mathcal{S}_{i,X}(\sigma^{-1} y)$. This formula defines a π -linear map $t' : B_k((N^*X)^{\otimes p}) \rightarrow \text{Hom}(W_{<i+1}, N^*X)$. It remains to check that $D't' = t' \cup t'$. Let e_i be any generator of W_i and $y \in FX$,

$$\begin{aligned} (t' \cup t')(y)(e_i) &= \mathcal{T}_{i,X}(s^i y) + (-1)^{|y|} t'(y)(\partial e_i) + t'(d_2 y)(e_i) \\ &= d\mathcal{S}_{i,X}(s^i y) - (-1)^{|\mathcal{S}_{i,X}|} \mathcal{S}_{i,X}(s^i d_1 y) + (-1)^{|y|} t'(y)(\partial e_i) + t'(d_2 y)(e_i) \\ &= d(t'(y)(e_i)) + t'(d_1 + d_2)(y)(e_i) + (-1)^{|y|} t'(y)(\partial e_i) \\ &= (D't')(y)(e_i). \end{aligned}$$

We have thus constructed a twisting cochain $t'_X \in \text{Hom}^1(B((N^*X)^{\otimes p}), \text{Hom}(W, N^*X))$ such that t'_X is natural in X and t'_X is π -linear.

2.3 Proof of part 2. Recall the notation introduced in [2]-Part II-§3. In particular, consider the simplicial model K of S^1 , and the cosimplicial space $\underline{X}$ defined as $\underline{X}(n) = \text{Map}(K(n), X)$ whose geometric realization $||\underline{X}||$ is homemorphic to X^{S^1} . Moreover, there is a natural equivalence $\psi_X : \text{Tot} C^* \underline{X} \rightarrow N^* ||\underline{X}||$, where $\text{Tot} C^* \underline{X}$ denotes the total complex of the simplicial complex $C^* \underline{X}$. On the other hand, J. Jones, [10]-6.3, has proved that there is natural chain map $\theta_X : \mathfrak{C} N^* X \rightarrow \text{Tot} C^* \underline{X}$ such that, if X is 1-connected, the composite $J_X = \psi_X \circ \theta_X$ induces an isomorphism $HH_* N^* X \cong H^* X^{S^1}$. We obtain the following diagram in the category of π -differential graded modules:

$$\begin{array}{ccc} \mathfrak{C}(\Omega B(N^*X)^{\otimes p}) & \xrightarrow{\phi \circ \mathfrak{C}\tilde{\kappa}_X} & \text{Hom}(W, \mathfrak{C} N^* X) \\ \mathfrak{C}(\alpha_{(N^*X)^{\otimes p}}) \downarrow & & \downarrow \text{Hom}(W, \theta_X \circ \psi_X) \\ \mathfrak{C}((N^*X)^{\otimes p}) & & \text{Hom}(W, N^* ||\underline{X}||) \\ sh^{(p)} \uparrow & & \uparrow \tilde{\Gamma}_X \\ (\mathfrak{C} N^* X)^{\otimes p} & \xrightarrow{\psi_X^{\otimes p}} & (N^* ||\underline{X}||)^{\otimes p} \end{array}$$

where Γ_X denotes any natural structural map defining the usual Steenrod operations, [13]. The functor $X \mapsto \mathfrak{C}((N^*X)^{\otimes p})$ preserves the units and is acyclic and the functor $X \mapsto \text{Hom}(W, N^* ||\underline{X}||)$ is corepresentable on the models $\mathcal{M}$, (see [2]-II-lemma 3.9). Therefore, by Corollary B.4, there exists a π -linear natural transformation $T_X : \mathfrak{C}((N^*X)^{\otimes p}) \rightarrow \text{Hom}(W, N^* ||\underline{X}||)$ such that $T_X \circ \mathfrak{C}\alpha_{(N^*X)^{\otimes p}} \simeq_\pi \text{Hom}(W, \theta_X \circ \psi_X) \circ \phi \circ \mathfrak{C}\tilde{\kappa}_X$ and $T_X \circ sh^{(p)} \simeq_\pi$

$\tilde{\Gamma}_X \circ \psi_X^{\otimes p}$. Consequently, the second part of theorem B is proved.

§3 π -shc models.

We continue the study of the homotopy category of π -shc algebras by constructing a convenient model of these algebras. As it would be expected this model is rather complicated. Nevertheless, in the topological case, it provides a theoretical object simpler than the singular cochain algebra.

3.1 Minimal algebra. Let $V = \{V^i\}_{i \geq 1}$ be a graded vector space and let TV denotes the free graded algebra generated by V : $T^r V = V \otimes V \otimes \dots \otimes V$ (r -times) and $v_1 v_2 \dots v_k \in (TV)^n$ if $\sum_{i=1}^k |v_i| = n$. The differential d on TV is the unique degree 1 derivation on TV defined by a given linear map $V \rightarrow TV$ and such that $d_V \circ d_V = 0$. The differential $d_V : TV \rightarrow TV$ decomposes as $d = d_0 + d_1 + \dots$ with $d_k V \subset T^{k+1} V$. In particular (V, d_0) is a differential graded vector space. If we assume that $V^1 = 0$ and $d_0 = 0$ then (TV, d_V) is called a *1-connected minimal algebra*. For any differential graded algebra (TU, d_U) such that $H^0(TU, d_U) = \mathbb{F}_p$ and $H^1(TU, d_U) = 0$ there exists a sequence of homomorphisms of differential graded algebras,

$$(TU, d_U) \xrightarrow{p_V} (TV, d_V) \xrightarrow{\varphi_V} (TU, d_U)$$

where (TV, d_V) denotes a 1-connected minimal algebra, $p_V \circ \varphi_V = id$, $\varphi_V \circ p_V \simeq_{\mathbf{DA}} id$ and V such that $V \cong H(U, d_{U,0})$. Moreover, [8], [6], [7], (TV, d_V) is unique up to isomorphisms.

3.2 Minimal model of a product. Assume that (A, d_A) is a differential graded algebra such that $H^0(A, d_A) = \mathbb{F}_p$ and $H^1(A, d_A) = 0$, and let $(TU[n], d_{U[n]}) = \Omega((BA)^{\otimes n})$, $n \geq 1$. By the discussion above, for each $n \geq 1$, we obtain a sequences:

$$(TU[n], d_{U[n]}) = \Omega((BA)^{\otimes n}) \xrightarrow{p_{V[n]}} (TV[n], d_{V[n]}) \xrightarrow{\varphi_{V[n]}} (TU[n], d_{U[n]})$$

$$\text{with } V[n] = s^{-1} \left(\overline{H((BA)^{\otimes n})} \right) \cong s^{-1} \left(\overline{(H(BA))^{\otimes n}} \right) = s^{-1} \left(\overline{(\mathbb{F}_p \oplus sV)^{\otimes n}} \right) \\ \cong \left(\bigoplus_{k=1}^n \left((\mathbb{F}_p)^{\otimes k-1} \otimes V \otimes (\mathbb{F}_p)^{\otimes n-k} \right) \right) \oplus \dots \oplus s^{-1} (sV \otimes sV \otimes \dots \otimes sV).$$

For $n = 1$, $V[1] = V = s^{-1} \overline{H(BA)}$ and the composite $\psi_V = \alpha_A \circ \varphi_V : (TV, d_V) \rightarrow A$ is a quasi-isomorphism. The algebra (TV, d_V) is called the *1-connected minimal model* of A .

For $n \geq 2$, consider the homomorphism $q_{V[n]} : (TV[n], d_{V[n]}) \rightarrow (TV, d_V)^{\otimes n}$ defined by $q_{V[n]}(y) = 1^{\otimes k-1} \otimes y \otimes 1^{\otimes n-k}$ if $y \in V_k := \mathbb{F}_p^{\otimes k-1} \otimes V \otimes \mathbb{F}_p^{\otimes n-k}$, $k = 1, 2, \dots, n$ et $q_{V[n]}(y) = 0$ if $y \in V[n] - \bigoplus_{i=1}^n V_i$. The composite $(TV[n], d_{V[n]}) \xrightarrow{q_{V[n]}} (TV, d_V)^{\otimes n} \xrightarrow{(\psi_V)^{\otimes n}} (A, d_A)^{\otimes n}$ is a quasi-isomorphism ([2]-6.5). Therefore $(TV[n], d_{V[n]})$ is the minimal model of $A^{\otimes n}$.

3.3 Lemma. (Equivariant lifting lemma.) *Let π any finite group and consider*

- i) *a graded free π -module, $U = \{U^i\}_{i \geq 2}$,*
- ii) *a minimal model, (TU, d) , with π acting by the rule $\sigma \cdot u_1 u_2 \dots u_k = \sigma u_1 \dots \sigma u_k$, so that (TU, d) is a π -complex,*
- iii) *two differential graded algebras, A and B , which are also π -complexes,*
- iv) *a homomorphism of differential graded algebras, $f : (TU, d) \rightarrow B$, which is π -equivariant,*
- v) *a surjective quasi-isomorphism, $\varphi : A \rightarrow B$, of differential graded algebras which is π -equivariant.*

Then there exists a homomorphism of differential graded algebras $g : (TU, d) \rightarrow B$ which is π -equivariant and such that $\varphi \circ g = f$.

Proof. We choose a homogeneous linear basis $\{u_i\}_{i \in I}$ of the π -free graded module U with I a well-ordered set. We denote by $U_{<i}$ the sub- π -module generated by the elements u_i , $j < i$. The minimality condition together with the 1-connectivity condition imply that $du_i \in T(U_{<i})$. Suppose that $g : (T(U_{<i}), d) \rightarrow B$ has been constructed so that g is π -equivariant and $\varphi \circ g = f$. Then $g(du_i) \in B \cap \text{Ker } d_B$ and $\varphi \circ g(du_i) = d_A f(e_i)$. Therefore there exists $b_i \in B$ such that $g(du_i) = d_B b_i$. Moreover, $d_A f(u_i) = d_A \varphi b_i$. For $\sigma \in \pi$, we set $g(\sigma u_i) = \sigma(b_i + z_i)$, and we check that $\varphi \circ g(\sigma u_i) = f(\sigma u_i)$ and $dg(\sigma u_i) = g(d\sigma u_i)$.

3.4 π -shc-minimal model. For any integer $n > 1$ the group $\mathfrak{S}_n$ acts on $V[n] \subset s^{-1}(H(BA))^{\otimes n}$. This action extends diagonally on $TV[n]$ so that the differential $d_{V[n]}$ and the homomorphism $(\psi_V)^{\otimes n} \circ q_{V[n]}$ are $\mathfrak{S}_n$ -linear. Since the natural map $\alpha_{A^{\otimes n}}$ is a $\mathfrak{S}_n$ -equivariant surjective quasi-isomorphism, Lemma 3.3 implies that the composite $(\psi_V)^{\otimes p} \circ q_{V[n]}$ lifts to a homomorphism of differential graded algebras $L : TV[n] \rightarrow \Omega B(A^{\otimes n})$ which is $\mathfrak{S}_n$ -equivariant and $\alpha_{A^{\otimes n}} \circ L = (\psi_V)^{\otimes n} \circ q_{V[n]}$. Let (A, d_A, μ_A) be an augmented shc algebra and assume that $H^0(A, d_A) = \mathbb{F}_p$ and that $H^1(A, d_A) = 0$. The composite, $\mu_V^{(n)} := p_V \circ \mu_A^{(n)} \circ L : (TV[n], d_{V[n]}) \rightarrow (T(V), d_V)$ is a homomorphism of differential graded algebras. If $n = 2$, $\mu_V^{(2)} := \mu_V : (T(V[2]), d_{V[2]}) \rightarrow (TV, d)$. The triple (TV, d_V, μ_V) is called a *shc-minimal model* for (A, d_A, μ_A) (See [2]-I-§6 for more details.)

Let $(A, d_A, \mu_A, \tilde{\kappa}_A)$ be a π -shc algebra such that $H^0 A = \mathbb{F}_p$ and $H^1 A = 0$. By Lemma 3.3, $\tilde{\kappa}_A \circ L$ lifts to a π -equivariant homomorphism of algebras $\hat{\kappa}_A : TV[p] \rightarrow \text{Hom}(W, \Omega BA)$. Since π is assumed to act trivially on TV the composite $\tilde{\kappa}_V = \text{Hom}(W, p_V) \circ \hat{\kappa}_A : TV[p] \rightarrow \text{Hom}(W, TV)$ is a π -equivariant map and we obtain the commutative diagram:

$$\begin{array}{ccc} \Omega B(A^{\otimes p}) & \xrightarrow{\tilde{\kappa}_A} & \text{Hom}(W, A) \\ \uparrow L & & \uparrow \text{Hom}(W, \alpha_A) \\ (TV[p], d_{V[p]}) & \xrightarrow{\hat{\kappa}_A} & \text{Hom}(W, \Omega BA) \\ & \searrow \tilde{\kappa}_V & \downarrow \text{Hom}(W, p_V) \\ & & \text{Hom}(W, TV). \end{array}$$

The triple $(TV, d_V, \mu_V, \tilde{\kappa}_V)$ is called a *π -shc minimal model* for the π -shc algebra $(A, d_A, \mu_A, \tilde{\kappa}_A)$.

3.5. Proposition. Let $(A, d_A, \mu_A, \tilde{\kappa}_A)$ and $(TV, d_V, \mu_V, \tilde{\kappa}_V)$ be as above then:

- 1) $(TV, d_V, \mu_V, \tilde{\kappa}_V)$ is a π -shc algebra,
- 2) $\psi_V : (TV, d) \rightarrow A$ is a strict π -shc homomorphism and a quasi-isomorphism,
- 3) the canonical maps $(TV, d_V) \xrightarrow{\iota_V} \mathfrak{C}(TV, d_V) \xrightarrow{\rho_V} B(TV, d_V)$ respect the Steenrod operation in homology.

- 4) the quasi-isomorphism $\psi_V : (TV, d_V) \rightarrow (A, d_A)$ induces a commutative diagram

$$\begin{array}{ccccc} (TV, d_V) & \xrightarrow{\iota_V} & \mathfrak{C}(TV, d_V) & \xrightarrow{\rho_V} & B(TV, d_V) \\ \psi_V \downarrow & & \downarrow \mathfrak{C}\psi_V & & \downarrow B\psi_V \\ (A, d_A) & \xrightarrow{\iota_A} & \mathfrak{C}(A, d_A) & \xrightarrow{\rho_A} & B(A, d_A) \end{array}$$

in which all vertical arrows induce isomomorphisms in homology which respect the Steenrod operations.

Proof. From the definition of $\tilde{\kappa}_V$, we deduce that $ev_0 \circ \hat{\kappa}_V \simeq_{\mathbf{DA}} \mu_V^n$. This implies that $(TV, \mu_V, \tilde{\kappa}_V)$ is a π -shc algebra. Since $p_V \circ \varphi_V = id_{TV}$, $\varphi_V \circ p_V \simeq_{\mathbf{DA}} id_{\Omega BA}$ then $Hom(W, \phi_V) \circ \tilde{\kappa}_V \simeq_{\pi-\mathbf{DA}} \hat{\kappa}_A$. Indeed, it is straightforward to check that if π acts trivially on A and A' and if $f, g : A \rightarrow A'$ are homotopic in $\mathbf{DA}$, then $Hom(W, f)$ and $Hom(W, g)$ are π -homotopic in $\mathbf{DA}$. It remains to show that ψ_V is a homomorphism π -shc algebras. This follows directly from the commutativity up to π -homotopy in $\mathbf{DA}$ of the diagram

$$\begin{array}{ccc} \Omega((BA)^{\otimes p}) & \xrightarrow{\tilde{\kappa}_A} & Hom(W, A) \\ L \uparrow & & \uparrow Hom(W, \psi_V) \\ TV[p] & \xrightarrow{\tilde{\kappa}_V} & Hom(W, TV). \end{array}$$

3.6. Proposition. Let $(A, d_A, \mu_A, \tilde{\kappa}_A)$ and $(TV, d_V, \mu_V, \tilde{\kappa}_V)$ be as above. Assume $p = 2$ and set $\hat{V} = V[2]$, $V' = V \otimes \mathbb{F}_p$, $V'' = \mathbb{F}_p \otimes V$ and $V' \# V'' = s^{-1}(sV \otimes sV)$, so that: $\hat{V} = V' \oplus V'' \oplus V' \# V''$. If $\tilde{\kappa} : \Omega B(A \otimes A) \rightarrow Hom(W, A)$ satisfies:

$$(\star) \quad \tilde{\kappa}(\langle [a \otimes b] \rangle) = 0 \text{ if } [a \otimes b] \in B_1(A \otimes A) \text{ and } \epsilon_A(a) = 0 \text{ or } \epsilon_A(b) = 0,$$

then, for any $v \in V^n \cap \ker d_V$, $Sq^i(cl(v)) = \tilde{\kappa}_V(v \# v)(e_{n-i-1})$.

Observe that if $A = N^*X$, condition $(\star)$ is satisfied (see 2.2, $(\star_1)$).

Proof. Define $K_A : \Omega(BA \otimes BA) \rightarrow \Omega BA \otimes \Omega BA$ as follows: we write $\Omega(BA \otimes BA) = (T(s^{-1}(B^+A \otimes \mathbb{F}_p) \oplus s^{-1}(\mathbb{F}_p \otimes B^+A) \oplus s^{-1}(B^+A \otimes B^+A)), D)$ and we set $K_A(s^{-1}[a_1 | \dots | a_i] \otimes 1) = s^{-1}[a_1 | \dots | a_i] \otimes 1$, $K_A(1 \otimes s^{-1}[a_1 | \dots | a_i]) = 1 \otimes s^{-1}[a_1 | \dots | a_i]$, $K_A(s^{-1}([a_1 | \dots | a_i] \otimes [b_1 | \dots | b_j])) = 0, i, j > 0$. One easily check ([2]-I-6.4) that K_A commutes with the differentials. Since, $(\alpha_A \otimes \alpha_A) \circ K_A = \alpha_{A \otimes A} \circ \Omega \overline{sh}$, the homomorphism K_A is a surjective quasi-isomorphism. The map $S : W \otimes_{\pi} (A \otimes A) \rightarrow W \otimes_{\pi} \Omega(BA \otimes BA)$ defined by $S(e_0 \otimes a \otimes b) = e_0 \otimes \langle [a] \otimes 1 \rangle \langle 1 \otimes [b] \rangle$ and if $i > 0$ by $S(e_i \otimes a \otimes b) = e_i \otimes \langle [a] \otimes 1 \rangle \langle 1 \otimes [b] \rangle + e_{i-1} \otimes \langle [b] \otimes [a] \rangle$ satisfies: $id_W \otimes (\alpha_A \otimes \alpha_A) \circ id_W \otimes K_A \circ S = id_{W \otimes_{\pi} A \otimes A}$ and $Sd = dS$. As in 1.6, we deduce:

$$\begin{aligned} Sq^i x &= cl(\tilde{\kappa}(i_{A \otimes A}(a \otimes a))(e_{n-i})) = cl(\kappa(e_{n-i} \otimes i_{A \otimes A}(a \otimes a))) = cl(\kappa_{/\pi}(e_{n-i} \otimes i_{A \otimes A}(a \otimes a))) \\ &= cl(\kappa_{/\pi}(((id \otimes \Omega \overline{sh}) \circ S)(e_{n-i} \otimes i_{A \otimes A}(a \otimes a)))) \\ &= cl(\kappa_{/\pi}(e_{n-i} \otimes \langle [a \otimes 1] \rangle \langle [1 \otimes a] \rangle)) + cl(\kappa_{/\pi}(e_{n-i-1} \otimes \Omega \overline{sh}(\langle [a] \otimes [a] \rangle))) \\ &= cl(\tilde{\kappa}(\Omega \overline{sh}(\langle [a] \otimes [a] \rangle))(e_{n-i-1})). \end{aligned}$$

The last equality comes from the condition $(\star)$ and the fact that $\Omega \overline{sh}$ is π -linear.

Now observe that $\varphi_{\hat{V}} : T(V' \oplus V'' \oplus V' \# V'') \rightarrow T(s^{-1} \overline{BA} \otimes \overline{BA})$ identifies $v' \in V'$ with $[a] \otimes 1$, $v'' \in V''$ with $1 \otimes [a]$ and $v' \# v'' \in V' \# V''$ with $\langle [a] \otimes [a] \rangle$. Thus if $v \in V^n \cap \ker d_V$ and $x = cl(v)$ then $H\psi_V \circ Sq^i(cl(v)) = cl(\tilde{\kappa}(\Omega \overline{sh} \circ \varphi_{\hat{V}}(v' \# v'')))$. The formula now follows from the definition of $\tilde{\kappa}_V$.

3.7 shc-equivalence. Two shc algebras (resp. π -shc algebras) A and A' are shc equivalent, $A \simeq_{shc} A'$ (resp. π -shc equivalent, $A \simeq_{\pi-shc} A'$) if there exists a sequence of strict shc (resp. π -shc) homomorphism $A \leftarrow A_1 \rightarrow \dots \rightarrow A'$ inducing isomorphisms in homology. If $A \simeq_{\pi-shc} A'$, then $H(A) \cong H(A')$, and A, A' have the same π -shc minimal model. Two spaces X and Y are shc equivalent (resp. π -shc equivalent) if the differential graded algebras N^*X and N^*Y are shc equivalent (resp. π -shc equivalent).

3.8. Example. We exhibit two spaces X and Y which are shc equivalent but not π -shc equivalent. Let us consider $X = \Sigma^2 \mathbb{C}P^2$ and $Y = S^4 \vee S^6$. The spaces X and Y have the same Adams-Hilton model namely $(T(x_3, x_5), 0)$. This shows, in particular, that $H_* \Omega X \cong H_* \Omega Y$, as graded Hopf algebras, and that X and Y are $\mathbb{F}_2$ -formal.

Furthermore, $H^* = H^*X = \mathbb{F}_2 a_4 \oplus \mathbb{F}_2 a_6 \cong H^*Y$ as graded algebras (with trivial products). Hence X and Y have the same minimal model say (TV, d_V) with $V = s^{-1}H_+\Omega X = s^{-1}T^+(x_3, x_5)$, $d_V s^{-1}x_3 = d_V s^{-1}x_5 = 0$. The map $\psi : (TV, d_V) \rightarrow H^*$ defined by $\psi(s^{-1}x_3) = a_4$, $\psi(s^{-1}x_5) = a_6$ and $\psi = 0$ on $s^{-1}T^{\geq 2}(x_3, x_5)$ is a surjective quasi-isomorphism. Let $(TV, d_V, \mu_1, \kappa_1)$ (resp. $(TV, d_V, \mu_2, \kappa_2)$) be a π -shc minimal model of X (resp. Y), then for $i = 1, 2$ $\mu_i : (T\hat{V}, d_{\hat{V}}) \rightarrow (TV, d_V)$ identifies the two copies of V . Moreover, $|\mu_i(x\#y)| > 6$ for any $x, y \in V$. Therefore $\psi \circ \mu_i(x\#y) = 0$. This yields the commutative diagram

$$\begin{array}{ccccc} T\hat{V} & \xrightarrow{\psi \otimes \psi \circ q_{\hat{V}}} & H^* \otimes H^* & \xleftarrow{\psi \otimes \psi \circ q_{\hat{V}}} & T\hat{V} \\ \mu_1 \downarrow & & \downarrow m_{H^*} & & \downarrow \mu_2 \\ TV & \xrightarrow{\psi} & H^* & \xleftarrow{\psi} & TV \end{array}$$

This proves that X and Y are shc equivalent. Now we prove that X and Y are not π -shc equivalent. Recall that if $a_4 \in H^4 \Sigma^2 \mathbb{C}P^2$ then $Sq^2 a_4 = a_6$ while if $a_4 \in H^4(S^4 \vee S^6)$ then $Sq^2 a_4 = 0$. Thus, by Proposition 3.6, $\tilde{\kappa}_1(a_4 \# a_4)(e_1) = a_6$ while $\tilde{\kappa}_2(a_4 \# a_4)(e_1) = 0$. Now $\tilde{\kappa}_1, \tilde{\kappa}_2$ are not π -homotopic, and hence X and Y are not π -shc equivalent.

3.9. Example. We compute the Steenrod operations on HH_*A when $A = \mathbb{F}_2[u] = T(u)$, $|u| = 2$.

First, recall that, endowed with the shuffle product, $\mathfrak{C}T(u) \xrightarrow{sh} \mathfrak{C}(T(u) \otimes T(u)) \xrightarrow{\mathfrak{C}m} \mathfrak{C}T(u)$, the complex $\mathfrak{C}T(u)$ is a commutative differential graded algebra. We consider the homomorphism of differential graded algebras $\rho : (T(u) \otimes \Lambda su, 0) \rightarrow \mathfrak{C}T(u)$ defined by $\rho(u \otimes 1) = u[\]$, $\rho(1 \otimes su) = 1[u]$. As proved in [11]-Proposition 3.1.2, ρ is a quasi-isomorphism of differential graded algebras. In particular, $T(u) \otimes \Lambda su \cong HH_*T(u)$ as commutative graded algebras.

Secondly, we define a structure of π -shc algebra on $T(u)$. Let $V = u\mathbb{F}_2$ and set $\hat{V} = u'\mathbb{F}_2 \oplus u''\mathbb{F}_2 \oplus u' \# u''\mathbb{F}_2$. We define $\tilde{\kappa} : T(\hat{V}) \rightarrow \text{Hom}(W, T(u))$ by $\tilde{\kappa}(u')(e_i) = \tilde{\kappa}(u')(e_i \tau) = \tilde{\kappa}(u'')(e_i) = \tilde{\kappa}(u'')(e_i \tau) = u$ if $i = 0$ and $= 0$ if $i > 0$ and $\tilde{\kappa}(u' \# u'')(e_i) = \tilde{\kappa}(u' \# u'')(e_i \tau) = u$ if $i = 1$, $= 0$ if $i > 0$. As in the proof of Proposition 3.6, $\tilde{\kappa}$ defines the Steenrod operations on $HT(u) = T(u)$: $Sq^0(u) = u$, $Sq^2 = u^2$ and $Sq^i u = 0$ if $i \neq 0, 2$. By Proposition 3.6 one deduces that $\tilde{\kappa}$ is the unique π -shc structural map on $T(u)$ with the Steenrod operations defined above.

Now in order to compute the Steenrod operations on $HH_*T(u)$ we consider the structural map $\tilde{\theta} = \phi \circ \mathfrak{C}\tilde{\kappa} : \mathfrak{C}(T\hat{V}) \rightarrow \text{Hom}(W, \mathfrak{C}T(u))$, and a linear section S of $id_W \otimes \mathfrak{C}q_{\hat{V}} : W \otimes_{\pi} \mathfrak{C}(T\hat{V}) \rightarrow W \otimes_{\pi} \mathfrak{C}(T(u) \otimes T(u))$. Then Steenrod operations are defined by: $Sq^i(x) = cl(\theta(S \circ (id_W \otimes sh)(e_{n-i} \otimes x \otimes x))$, $x \in HH^n T(u)$.

Finally observe that S is uniquely determined in low degrees as follows:

$$\begin{array}{llll} S(e_i \otimes 1[\]) & = e_i \otimes 1[\] & S(e_i \otimes (u \otimes 1)[\]) & = e_i \otimes u'[\] \\ S(e_i \otimes (1 \otimes u)[\]) & = e_i \otimes u''[\] & S(e_i \otimes 1[u \otimes 1]) & = e_i \otimes 1[u'] \\ S(e_i \otimes (u \otimes u)[\]) & = e_i \otimes u'u''[\] + e_{i-1} \otimes u' \# u''[\] & S(e_i \otimes 1[1 \otimes u]) & = e_i \otimes 1[u''] \\ S(e_i \otimes (u \otimes 1)[u \otimes 1]) & = e_i \otimes u'[u'] & S(e_i \otimes (1 \otimes u)[1 \otimes u]) & = e_i \otimes u''[u''] \\ S(e_i \otimes (1 \otimes u)[u \otimes 1]) & = e_i \otimes u''[u'] + e_i \otimes u' \# u''[\] & & \\ S(e_i \otimes (u \otimes 1)[1 \otimes u]) & = e_i \otimes u'[u''] + e_i \otimes u' \# u''[\] & & \\ S(e_i \otimes 1[u \otimes u]) & = e_i \otimes 1[u'u''] + e_{i-1} \otimes 1[u' \# u''] & S(e_i \otimes 1[u \otimes 1][1 \otimes u]) & = e_i \otimes 1[u'|u''] \\ S(e_i \otimes 1[1 \otimes u][u \otimes 1]) & = e_i \otimes 1[u''|u'] + e_i \otimes 1[u' \# u''] & & \end{array}$$

(Here we make the convention that $e_{-1} = 0$.) Therefore, since $\tilde{\theta}$ respects the products

and with the aid of Proposition 3.6, we can do the following computations:

$$\begin{aligned}
Sq^0(u[]) &= cl(\theta \circ S \circ sh(e_2 \otimes u[] \otimes u[])) = cl(\theta(e_2 \otimes u'u''[] + e_1 \otimes u' \# u''[])) \\
&= cl(\tilde{\theta}(u'u'')(e_2)[]) + cl(\tilde{\theta}((u' \# u'')(e_1)[])) = cl(\tilde{\kappa}(u' \# u'')(e_1)[]) = Sq^0(u[]) = u[] \\
Sq^1(u[]) &= cl(\theta \circ S \circ sh(e_1 \otimes u[] \otimes u[])) = cl(\theta(e_1 \otimes u'u''[] + e_0 \otimes 1[u' \# u''])) \\
&= cl(\tilde{\kappa}(u' \# u'')(e_0)[]) = 0 \\
Sq^2(u[]) &= cl(\theta \circ S \circ sh(e_0 \otimes u[] \otimes u[])) = cl(\tilde{\kappa}(u'u'')(e_0)[]) = u^2. \\
Sq^0(1[u]) &= cl(\theta \circ S \circ sh(e_1 \otimes 1[u] \otimes 1[u])) = cl(\theta(e_1 \otimes 1[u'u''] + e_1 \otimes 1[u' \# u''])) \\
&= cl(1[\tilde{\kappa}(u' \# u'')(e_1)]) = 1[Sq^0 u] = 1[u] \\
Sq^1(1[u]) &= cl(\theta \circ S \circ sh(e_0 \otimes 1[u] \otimes 1[u])) \\
&= cl(\theta(e_0 \otimes 1[u'u''] + e_0 \otimes 1[u''|u'] + e_0 \otimes 1[u' \# u''])) \\
&= cl(1[u|u] + 1[u|u]) + cl(1[\tilde{\kappa}(u' \# u'')(e_1)]) = 0.
\end{aligned}$$

Thus the Steenrod operations are completely defined on $HH_*T(u) = T(u) \otimes \Lambda su$, by the Cartan formula and the formulas: $Sq^0(u[]) = u[]$, $Sq^2(u[]) = u^2[]$, $Sq^i(u[]) = 0$ for $i \neq 1, 2$, $Sq^0(1[u]) = 1[u]$, $Sq^i(1[u]) = 0$ if $i > 0$. We recover the topological Steenrod squares on $H^*((\mathbb{C}P^\infty)^{S^1}) = H^*(\mathbb{C}P^\infty) \otimes H^*(S^1)$. The technics developped in this example, can be performed in order to study the case when $A = \mathbb{F}_2[u]/(u^k)$, $k \geq 2$.

Appendix A - Technicalities on *shc*-algebras.

In this appendix we lay the material necessary in order to complete the proof of Theorem A.

A.1 Lemma. *Let A, A' be two *shc* algebras and let $f, g \in \mathbf{DA}(A, A')$ be such that $f \simeq_{\mathbf{DA}} g$. If f is a *shc* homomorphism then g is a *shc* homomorphism.*

Proof Let θ be a (f, g) -derivation such that $f - g = d_{A'} \circ \theta + \theta \circ d_A$. Then θ defines a coderivation $\theta' : BA \rightarrow BA'$: $\theta'([a_1] \dots [a_k]) = \sum_{i=1}^k (-1)^{\epsilon_i} [f a_1 | \dots | f a_{i-1} | \theta a_i | \dots | g a_{i+1} | \dots | g a_k]$ where $\epsilon_i = \sum_{j=1}^{i-1} |a_j| + (i-1)$ satisfying $Bf - Bg = d_{BA'} \circ \theta' + \theta' \circ d_{BA}$. Now let $f_1, f_2 \in \mathbf{DC}(C, C')$ and consider the adjoint homomorphism $\tilde{f}_i : C \rightarrow C' \rightarrow B\Omega C'$ ($i = 1, 2$). If we assume that $f_1 \simeq_{\mathbf{DC}} f_2$ then $\tilde{f}_1 \simeq_{\mathbf{DC}} \tilde{f}_2$ and thus, by [15]-1.11, $\Omega f_1 \simeq_{\mathbf{DA}} \Omega f_2$. Therefore if $f \simeq_{\mathbf{DA}} g$ then $\Omega Bf \simeq_{\mathbf{DA}} \Omega Bg$. The lemma follows now, from the obvious relations: $\mu_{A'} \circ \Omega B(g \otimes g) \simeq_{\mathbf{DA}} \mu_{A'} \circ \Omega B(f \otimes f) \simeq_{\mathbf{DA}} \Omega Bf \circ \mu_A \simeq_{\mathbf{DA}} \Omega Bg \circ \mu_A$.

A.2 Trivialized extensions. Let $\mathbf{TEX}_A$ be the category of trivialized extensions of A in $\mathbf{DA}$ in the sense of [15]-2.1: $X \xrightarrow{\alpha} A \in \mathbf{TEX}$ if $\alpha \in \mathbf{DA}(X, A)$ and if there exist $\rho \in \mathbf{DM}(A, X)$, $h \in \mathbf{Hom}(X, X)$ such that, $\alpha \circ \rho = id_A$, $\rho \circ \alpha - id_X = d_X \circ h + h \circ d_X$, $\rho \circ \eta_A = \eta_X$, $\epsilon_X \circ \rho = \epsilon_A$, $\alpha \circ h = 0$, $h \circ \rho = 0$, $h \circ h = 0$. A (strict) morphism of trivialized extension, $f : (X \xrightarrow{\alpha} A, \rho, h) \rightarrow (X' \xrightarrow{\alpha'} A, \rho', h')$ is a homomorphism $f \in \mathbf{DA}(X, X')$ such that $f \circ \rho = \rho'$ and $f \circ h = h' \circ f$.

The following facts are proved in [15] or straightforward to prove.

F1 If $X \xrightarrow{f} A \in \mathbf{TEX}_A$ and $Y \xrightarrow{g} X \in \mathbf{TEX}_X$, then $Y \xrightarrow{f \circ g} A \in \mathbf{TEX}_A$.

F2 If $X \xrightarrow{f} A \in \mathbf{TEX}_A$ and $X' \xrightarrow{f'} A' \in \mathbf{TEX}_{A'}$, then $X \otimes X' \xrightarrow{f \otimes f'} A \otimes A' \in \mathbf{TEX}_{A \otimes A'}$.

F3 If A is a differential graded algebra then $\Omega BA \xrightarrow{\alpha_A} A$ is an initial object in $\mathbf{TEX}$.

F4 If (A, μ_A) and $(A', \mu_{A'})$ are *shc* algebras then there exists a natural homomorphism

$\mu_{A \otimes A'}$ such that $(A \otimes A', \mu_{A \otimes A'})$ is a *shc* algebra. In particular, $A^{\otimes n}, n \geq 2$ is naturally a *shc* algebra.

F5 The homomorphism $\alpha_{\Omega BA} : \Omega B \Omega BA \rightarrow \Omega BA$ admits a section $\theta_{\Omega BA} \in \mathbf{DA}(\Omega BA, \Omega B \Omega BA)$.

F6 If (A, μ_A) is a *shc* algebra so is ΩBA .

F7 $\Omega B \alpha_A \simeq_{\mathbf{DA}} \alpha_{\Omega BA}$.

F8 If (A, μ_A) is a *shc* algebra then $\alpha_A : \Omega BA \rightarrow A$ is a strict *shc* homomorphism.

A.3 Lemma. *Let (A, d, μ) be a *shc* algebra. For any $n \geq 2$, there exists a homomorphism of differential graded algebras (called the *shc iterated structural map*) $\mu^{(n)} : \Omega B(A^{\otimes n}) \rightarrow \Omega BA$ such that: $\mu^{(2)} = \mu$ and $\alpha_A \circ \mu^{(n)} \circ i_{A^{\otimes n}} \simeq m_A^{(n)}$. Moreover, $\mu^{(n)}$ is a strict homomorphism of *shc* algebras.*

Proof. Let A_1 and A_2 be two differential graded algebras. From F3, we deduce that there exists a natural homomorphism of differential graded algebras: $\theta_{A_1, A_2} : \Omega B(A_1 \otimes A_2) \rightarrow \Omega B(A_1 \otimes \Omega BA_2)$ such that: $(\star) \quad (id_{A_1} \otimes \alpha_{A_2}) \circ \alpha_{A_1 \otimes \Omega BA_2} \circ \theta_{A_1, A_2} = \alpha_{A_1 \otimes A_2}$. It follows, from the unicity property, that if A_3 is a differential graded algebra then $(\theta_{A_1, A_2} \otimes id_{\Omega BA_3}) \circ \theta_{A_1 \otimes A_2, A_3} = (id_{\Omega BA_1} \otimes \theta_{A_2, A_3}) \circ \theta_{A_1, A_2 \otimes A_3}$. Suppose inductively that $\mu^{(n)}$ is defined for some $n \geq 2$. We define $\mu^{(n+1)}$ as the composite $\Omega B(A^{\otimes n+1}) \xrightarrow{\theta_{A, A^{\otimes n}}} \Omega B(A \otimes \Omega B(A^{\otimes n})) \xrightarrow{\Omega B(id \otimes \mu^{(n)})} \Omega B(A \otimes \Omega B(A)) \xrightarrow{\Omega B(id \otimes \alpha_A)} \Omega B(A \otimes A) \xrightarrow{\mu} \Omega BA$. If we assume moreover that $\alpha_A \circ \mu^{(n)} \circ i_{A^{\otimes n}} = m_A^{(n)}$ then the identities $(\star)$ and $\alpha_- \circ i_- = id_-$ imply that $\alpha_A \circ \mu^{(n+1)} \circ i_{A^{\otimes n+1}} = m_A^{(n+1)}$. The last statement is Proposition 4.5 in [15].

A.4 Lemma. *Let (W, ψ_W) be the standard resolution.*

a) *If (A, d) is a differential graded algebra then $\text{Hom}(W, A) \xrightarrow{ev_0} A \in \mathbf{TEX}_A$.*

b) *If $X \xrightarrow{\alpha} A \in \mathbf{TEX}_A$ then $\text{Hom}(W, X) \xrightarrow{\text{Hom}(W, \alpha)} \text{Hom}(W, A) \in \mathbf{TEX}_{\text{Hom}(W, A)}$.*

c) *If A is a *shc* algebra then*

(i) *$\text{Hom}(W, A)$ is a *shc* algebra,*

(ii) *ev_0 is a *shc* homomorphism,*

(iii) *Let A' be a *shc* algebra. A homomorphism of differential graded algebras*

*$f : A' \rightarrow \text{Hom}(W, A)$ is a *shc* homomorphism if and only if $ev_0 \circ f$ is a *shc* homomorphism.*

Proof. a) Let us denote $[g_0, g_1, \dots, g_n] \in G^{\times n+1}$ a generator on W_n and recall the linear homotopy $k : W \rightarrow W$ defined by $k([g_0, \dots, g_n]) = \sum_{i=0}^n (-1)^i [g_0, \dots, g_{i-1}, 1, g_i, \dots, g_n]$. Obviously, $k \circ \partial_W + \partial_W \circ k = id_W - \eta_W \circ \epsilon_W$ and $k \circ k = 0$. We define $h : \text{Hom}(W, A) \rightarrow \text{Hom}(W, A)$ by $h(f) = k \circ f$. If $\rho : A \rightarrow \text{Hom}(W, A)$ is defined by $\rho(a)(w) = a \eta_A \epsilon_W(w)$, $a \in A$, $w \in W$, a straightforward computation shows that $h \circ D + D \circ h = id_{\text{Hom}(W, A)} - \rho \circ ev_0$ and $h \circ h = 0$. One also easily checks that $ev_0 \circ \rho = id_A$, $ev_0 \circ h = 0$, $h \circ \rho = 0$, $\rho \circ \eta_A = \eta_{\text{Hom}(W, A)}$, $\epsilon_{\text{Hom}(W, A)} \circ \rho = \epsilon_A$, where $\eta_{\text{Hom}(W, A)} = \eta_A \circ \epsilon_W$ and $\epsilon_{\text{Hom}(W, A)}(f) = \epsilon_A \circ f \circ \eta_W(1)$.

b) The proof is similar to the end of the proof of part a).

c) (i) Consider the differential linear map $\psi_A : \text{Hom}(W, A) \otimes \text{Hom}(W, A) \rightarrow \text{Hom}(W, A \otimes A)$ defined by $\psi_A(f \otimes g) = (f \otimes g) \circ \psi_W$. Then, $\text{Hom}(W, m_A) \circ \psi_A = \cup_{\text{Hom}(W, A)}$ and the following equalities prove that ψ_A is a homomorphism of differential graded algebras.

$$\begin{aligned}
\psi_A((f_1 \otimes g_1) \cup (f_2 \otimes g_2)) &= (-1)^{|f_2||g_1|} \psi_A((f_1 \cup f_2) \otimes (g_1 \cup g_2)) \\
&= (-1)^{|f_2||g_1|} ((f_1 \cup f_2) \otimes (g_1 \cup g_2)) \circ \psi_W \\
&= (-1)^{|f_2||g_1|} (m_A \circ (f_1 \otimes f_2) \circ \psi_W) \otimes m_A \circ ((g_1 \otimes g_2) \circ \psi_W) \circ \psi_W \\
&= m_{A \otimes A} \circ ((f_1 \otimes g_1) \otimes (f_2 \otimes g_2)) \circ \psi_W \circ \psi_W \\
&= m_{A \otimes A} \circ (\psi_A(f_1 \otimes g_1) \otimes \psi_A(f_2 \otimes g_2)) \circ \psi_W \\
&= \psi_A(f_1 \otimes g_1) \cup \psi_A(f_2 \otimes g_2).
\end{aligned}$$

We define $\mu_{\text{Hom}(W,A)} : \Omega B(\text{Hom}(W,A)^{\otimes 2}) \rightarrow \Omega B\text{Hom}(W,A)$ as the composite $\Omega B\text{Hom}(W, \alpha_A \circ \mu_A) \circ \theta_{\text{Hom}(W,A \otimes A)} \circ \Omega B\psi$ where $\theta_{\text{Hom}(W,A \otimes A)}$ denotes the unique homomorphism of differential graded algebras such that $\theta_{\text{Hom}(W,A \otimes A)} \circ \text{Hom}(W, \alpha_A \otimes \alpha_A) \circ \alpha_{\text{Hom}(W, \Omega B A \otimes \Omega B A)} = \alpha_{\text{Hom}(W, A \otimes A)}$. Existence and unicity of $\theta_{\text{Hom}(W,A \otimes A)}$ is a direct consequence of F3, F1, F2 and part b). It turns out that $(\text{Hom}(W,A), \mu_{\text{Hom}(W,A)})$ is a *shc* algebra.

c)-(ii). Consider for any A in $\mathbf{DA}$ the commutative diagram

$$\begin{array}{ccccc}
\Omega B A & \xleftarrow{\Omega B(ev_0)} & \Omega B\text{Hom}(W,A) & \xleftarrow{\alpha_{\Omega B\text{Hom}(W,A)}} & (\Omega B)^2 \text{Hom}(W,A) \\
& & \downarrow \alpha_{\text{Hom}(W,A)} & & \downarrow \Omega B \alpha_{\text{Hom}(W,A)} \\
\alpha_A \searrow & & \text{Hom}(W,A) & \xleftarrow{\alpha_{\text{Hom}(W,A)}} & \Omega B\text{Hom}(W,A) B \\
& & \downarrow ev_0 & & \downarrow \Omega B ev_0 \\
& & A & \xleftarrow{\alpha_A} & \Omega B A.
\end{array}$$

From F3, we deduce the existence of $\theta'_{\Omega B A} : \Omega B A \rightarrow \Omega B\text{Hom}(W,A)$ such that $ev_0 \circ \alpha_{\text{Hom}(W,A)} \circ \theta'_{\Omega B A} = \alpha_A$ and $\theta'_{\Omega B A} \circ \Omega B\text{Hom}(W, ev_0) \simeq_{\mathbf{DA}} \alpha_{\Omega B\text{Hom}(W,A)}$. By F7, $\Omega B \alpha_{\text{Hom}(W,A)} \simeq_{\mathbf{DA}} \alpha_{\Omega B\text{Hom}(W,A)}$ and by F5, there is a section for $\alpha_{\Omega B\text{Hom}(W,A)}$ so that $(\star_A) \quad \theta'_{\Omega B A} \circ \Omega B ev_0 \simeq_{\mathbf{DA}} id_{\Omega B\text{Hom}(W,A)}$. Moreover, from the unicity property in F3, we know that $\Omega B ev_0 \circ \theta'_{\Omega B A} = id_{\Omega B A}$. In the following diagram, the left hand square is commutative,

$$\begin{array}{ccccc}
\Omega B A & \xleftarrow{\Omega B(ev_0)} & \Omega B\text{Hom}(W,A) & \xleftarrow{\theta'_{\Omega B A}} & \Omega B A \\
\mu_A \uparrow & & \uparrow \Omega B\text{Hom}(W, \alpha_A \circ \mu_A) & & \uparrow \mu_A \\
\Omega B(A^{\otimes 2}) & \xleftarrow{\alpha_{\Omega B(A^{\otimes 2})} \circ \Omega B(ev_0)} & \Omega B\text{Hom}(W, \Omega B(A^{\otimes 2})) & \xleftarrow{\theta'_{\Omega B(A^{\otimes 2})}} & \Omega B(A^{\otimes 2}).
\end{array}$$

Since the composite of horizontal maps are the identity maps, relations $(\star_A)$ and $(\star_{A \otimes A})$ imply that the right hand square commutes up to homotopy in $\mathbf{DA}$. Therefore, in the following diagram the upper square commutes, up to homotopy in $\mathbf{DA}$.

$$\begin{array}{ccccccc}
\Omega B\text{Hom}(W,A) & \xrightarrow{\Omega B(ev_0)} & \Omega B A & \xrightarrow{\theta'_{\Omega B A}} & \Omega B\text{Hom}(W,A) & & \\
\parallel & & & & \uparrow \theta'_{\Omega B A} \circ \mu_A & & \\
\Omega B\text{Hom}(W,A) & \xrightarrow{\Omega B\text{Hom}(W, \alpha_A \circ \mu_A)} & \Omega B\text{Hom}(W, \Omega B(A^{\otimes 2})) & \xleftarrow{\theta'_{\Omega B(A^{\otimes 2})}} & \Omega B(A^{\otimes 2}) & & \\
\mu_{\text{Hom}(W,A)} \uparrow & & \Omega B\text{Hom}(W, \Omega B(A^{\otimes 2})) \downarrow & & \parallel & & \\
\Omega B(\text{Hom}(W,A)^{\otimes 2}) & \xrightarrow{\Omega B\psi_A} & \Omega B\text{Hom}(W, A^{\otimes 2}) & \xleftarrow{\theta'_{\Omega B(A^{\otimes 2})}} & \Omega B(A^{\otimes 2}) & & \\
\Omega B(ev_0 \otimes ev_0) \downarrow & & \Omega B(ev_0) \downarrow & & \parallel & & \\
\Omega B(A^{\otimes 2}) & = & \Omega B(A^{\otimes 2}) & = & \Omega B(A^{\otimes 2}) & &
\end{array}$$

It is not difficult, if tedious, to check that the other cells in the diagram commutes also up to homotopy in $\mathbf{DA}$. This shows that $\Omega B(ev_0) \circ \mu_{\text{Hom}(W,A)} \simeq_{\mathbf{DA}} \mu_A \circ \Omega B(ev_0 \otimes ev_0)$, i.e. ev_0 is a strict *shc* map.

c)-(iii) Let $f : A' \rightarrow \text{Hom}(W,A)$ be a homomorphism of differential graded algebras such that $\mu_A \circ \Omega(ev_0 \otimes ev_0) \circ \Omega B(f \otimes f) \simeq_{\mathbf{DA}} (\Omega B(ev_0 \circ f)) \circ \mu_{A'}$. Thus in the following diagram the bigger square commutes, up to homotopy in $\mathbf{DA}$.

$$\begin{array}{ccccc}
\Omega B A' & \xrightarrow{\Omega B f} & \Omega B \text{Hom}(W, A) & \xrightarrow{\Omega B ev_0} & \Omega B A \\
\mu_{A'} \uparrow & & \mu_{\text{Hom}(W, A)} \uparrow & & \uparrow \mu_A \\
\Omega B(A' \otimes A') & \xrightarrow{\Omega B(f \otimes f)} & \Omega B(\text{Hom}(W, A) \otimes \text{Hom}(W, A)) & \xrightarrow{\Omega B(ev_0 \otimes ev_0)} & \Omega B(A \otimes A).
\end{array}$$

The right hand square commutes, up to homotopy in $\mathbf{DA}$, by part c-(ii) and thus so does the left hand square. This proves that f is a strict *shc* homomorphism. Conversely, if f is a *shc* map, by part c-(ii), the composite $f \circ ev_0$ is also a strict *shc* homomorphism.

A.5 Lemma. *If $(A, d_A, \mu_A, \tilde{\kappa}_A)$ is a π -shc algebra then,*

- a) $\tilde{\kappa}$ is a strict homomorphism of *shc* algebras,
- b) $H^* \phi_A : H\mathfrak{C}(\text{Hom}(W, A)) \rightarrow H(\text{Hom}(W, \mathfrak{C}A))$ preserves the natural multiplications.

Proof. a) By lemma A.3, $\mu^{(p)}$ is a strict homomorphism of *shc* algebras. Since $ev_0 \circ \tilde{\kappa}_A \simeq_{\mathbf{DA}} \alpha_A \circ \mu_A^{(p)}$, from Lemma A.4-c)-(iii), we deduce that $\tilde{\kappa}_A$ is a strict *shc* homomorphism.

b) We obtain the two commutative diagrams (Diagrams A and B):

$$\begin{array}{ccccc}
(\mathfrak{C}\text{Hom}(W, A))^{\otimes 2} & \xrightarrow{\phi_A^{\otimes 2}} & \text{Hom}(W, \mathfrak{C}A)^{\otimes 2} & \xrightarrow{\psi_{\mathfrak{C}A}^{\otimes 2}} & \text{Hom}(W, (\mathfrak{C}A)^{\otimes 2}) \\
\downarrow sh & & & & \text{Hom}(W, sh) \downarrow \\
\mathfrak{C}(\text{Hom}(W, A)^{\otimes 2}) & \xrightarrow{\mathfrak{C}\psi_{\text{Hom}(W, A)}} & \mathfrak{C}\text{Hom}(W, A^{\otimes 2}) & \xrightarrow{\phi_{A^{\otimes 2}}} & \text{Hom}(W, \mathfrak{C}(A^{\otimes 2})) \\
\mathfrak{C}\alpha_{\text{Hom}(W, A)^{\otimes 2}} \uparrow & & \uparrow \mathfrak{C}\text{Hom}(W, \alpha_{A^{\otimes 2}}) & & \\
\mathfrak{C}\Omega B(\text{Hom}(W, A)^{\otimes 2}) & \xrightarrow{\mathfrak{C}\psi_{\Omega B(\text{Hom}(W, A)^{\otimes 2})}} & \mathfrak{C}\Omega B\text{Hom}(W, A^{\otimes 2}) & & ,
\end{array}$$

and

$$\begin{array}{ccc}
\mathfrak{C}\text{Hom}(W, A^{\otimes 2}) & \xrightarrow{\phi_{A^{\otimes 2}}} & \text{Hom}(W, \mathfrak{C}(A^{\otimes 2})) \\
\mathfrak{C}\text{Hom}(W, \alpha_{A^{\otimes 2}}) \uparrow & & \uparrow \text{Hom}(W, \mathfrak{C}\alpha_{A^{\otimes 2}}) \\
\mathfrak{C}\text{Hom}(W, \Omega B(A^{\otimes 2})) & \xrightarrow{\phi_{\Omega B(A^{\otimes 2})}} & \text{Hom}(W, \mathfrak{C}\Omega B(A^{\otimes 2})).
\end{array}$$

From, F3, Lemma A.4-b) and F8 we deduce the existence of the homomorphisms of differential graded algebras θ'_A , $\theta'_{A^{\otimes 2}}$, $\theta''_{A^{\otimes 2}}$ such that : $\text{Hom}(W, \alpha_A) \circ \theta'_A = \alpha_{\text{Hom}(W, A)}$, $\text{Hom}(W, \alpha_{A^{\otimes 2}}) \circ \alpha_{\text{Hom}(W, \Omega B(A^{\otimes 2}))} \circ \theta''_{A^{\otimes 2}} = \alpha_{\text{Hom}(W, A^{\otimes 2})}$ and $\text{Hom}(W, \alpha_{A^{\otimes 2}}) \circ \theta'_{A^{\otimes 2}} = \alpha_{\text{Hom}(W, A^{\otimes 2})}$. The desired commutativity of the diagrams C, and D below is clear from the construction of $\mu_{\text{Hom}(W, A)}$ (Proof of lemma A.4-c)-(i)):

$$\begin{array}{ccccc}
\mathfrak{C}\Omega B(\text{Hom}(W, A)^{\otimes 2}) & \xrightarrow{\mathfrak{C}\psi_{\Omega B(\text{Hom}(W, A)^{\otimes 2})}} & \mathfrak{C}\Omega B\text{Hom}(W, A^{\otimes 2}) & \xrightarrow{\mathfrak{C}\theta''_{A^{\otimes 2}}} & \mathfrak{C}\Omega B\text{Hom}(W, \Omega B(A^{\otimes 2})) \\
\mathfrak{C}\mu_{\text{Hom}(W, A)} \downarrow & & & & \mathfrak{C}\alpha_{\text{Hom}(W, \Omega BA)} \circ \downarrow \mathfrak{C}\Omega B\text{Hom}(W, \mathfrak{C}\mu_A) \\
\mathfrak{C}\Omega B\text{Hom}(W, A) & & \xrightarrow{\mathfrak{C}\theta'_A} & & \text{Hom}(W, \Omega BA),
\end{array}$$

and

$$\begin{array}{ccc}
\mathfrak{C}\Omega B\text{Hom}(W, A^{\otimes 2}) & \xrightarrow{\mathfrak{C}\theta'_{\Omega B(A^{\otimes 2})}} & \mathfrak{C}\text{Hom}(W, \Omega B(A^{\otimes 2})) \xrightarrow{\phi_{\Omega B(A^{\otimes 2})}} \text{Hom}(W, \mathfrak{C}\Omega B(A^{\otimes 2})) \\
\mathfrak{C}\theta''_{A^{\otimes 2}} \downarrow & \mathfrak{C}\alpha_{\text{Hom}(W, \Omega B(A^{\otimes 2}))} \nearrow & \downarrow \text{Hom}(W, \mathfrak{C}\mu_A) \\
\mathfrak{C}\Omega B\text{Hom}(W, \Omega B(A^{\otimes 2})) & & \\
\mathfrak{C}\alpha_{\text{Hom}(W, \Omega BA)} \circ \downarrow \mathfrak{C}\Omega B\text{Hom}(W, \mathfrak{C}\mu_A) & & \\
\mathfrak{C}\text{Hom}(W, \Omega BA) & \xrightarrow{\phi_{\Omega BA}} & \text{Hom}(W, \mathfrak{C}\Omega BA).
\end{array}$$

Now we choose linear sections of $\mathfrak{C}\alpha_{\text{Hom}(W, A^{\otimes 2})}$ (resp. of $\text{Hom}(W, \mathfrak{C}\alpha_{A^{\otimes 2}})$) and we define a product $m_{\mathfrak{C}\text{Hom}(W, A)}$ (resp. a cup product $m_{\text{Hom}(W, \mathfrak{C}A)}$) on $\mathfrak{C}\text{Hom}(W, A)$ (resp. on $\text{Hom}(W, \mathfrak{C}A)$). Then gluing together diagrams A, B, C, and D we deduce that the following diagram

$$\begin{array}{ccc}
(\mathfrak{C}\mathrm{Hom}(W, A))^{\otimes 2} & \xrightarrow{\phi_A^{\otimes 2}} & (\mathrm{Hom}(W, \mathfrak{C}A))^{\otimes 2} \\
m_{\mathfrak{C}\mathrm{Hom}(W, A)} \downarrow & & \downarrow m_{\mathrm{Hom}(W, \mathfrak{C}A)} \\
\mathfrak{C}\mathrm{Hom}(W, A) & \xrightarrow{\phi_A} & \mathrm{Hom}(W, \mathfrak{C}A)
\end{array}$$

commutes up to a linear homotopy.

A.6 Lemma. *Let (TV, d_V) be a differential graded algebra and assume that a finite group π acts freely on V . Let A be a π -differential graded algebra and $f, g \in \pi\text{-}\mathbf{DA}(TV, A)$. If $f \simeq_{\pi\text{-}\mathbf{DA}} g$ then $\mathfrak{C}f \simeq_{\pi} \mathfrak{C}g$.*

Proof. The closed model category framework provides a convenient language in which we prove the lemma. For our purpose we define the cylinder $I(TV, d)$ on (TV, d) :

$$\begin{array}{ccc}
(TV, d) & \searrow \partial_0 & \\
& I(TV, d) \xrightarrow{p} (TV, d) & \\
& \nearrow \partial_1 &
\end{array}$$

with $I(TV, d) := (T(V_0 \oplus V_1 \oplus sV), D)$, $\partial_0 V = V_0$, $\partial_1 V = V_1$, $pv_0 = pv_1 = v$, $psv = 0$, $D = d$ on V_0 and on V_1 , $Dsv = Sdv$ where S is the unique (∂_0, ∂_1) -derivation $S : TV \rightarrow T(V_0 \oplus V_1 \oplus sV)$ extending the graded isomorphism $s : V \rightarrow sV$. The free π -action on TV naturally extends to a free π -action on $I(TV, d)$ so that $I(TV, d)$ is a π -algebra and the maps p, ∂_0 and ∂_1 are π -equivariant quasi-isomorphisms. By definition $f \simeq_C g$ (resp. $f \simeq_{\pi\text{-}C} g$) if there exists $H \in \mathbf{DA}(I(TV, d), A)$ (resp. $H \in \pi\text{-}\mathbf{DA}(I(TV, d), A)$) such that $H\partial_0 = f$ and $H\partial_1 = g$. It is straightforward to check that $f \simeq_C g$ if and only if $f \simeq_{\mathbf{DA}} g$, [7], (resp. $f \simeq_{\pi\text{-}C} g$ if and only if $f \simeq_{\pi\text{-}\mathbf{DA}} g$). Consider the commutative diagram

$$\begin{array}{ccc}
\mathfrak{C}TV \oplus \mathfrak{C}TV & \xrightarrow{\mathfrak{C}id \oplus \mathfrak{C}id} & \mathfrak{C}TV \\
\mathfrak{C}\partial_0 \oplus \mathfrak{C}\partial_1 \downarrow & & \mathfrak{C}p \uparrow \\
\mathfrak{C}ITV & \xrightarrow{id} & \mathfrak{C}ITV.
\end{array}$$

Now $\mathfrak{C}\partial_0 \oplus \mathfrak{C}\partial_1$ is injective and its cokernel is a projective π -module. Hence, $\mathfrak{C}\partial_0 \oplus \mathfrak{C}\partial_1$ is a cofibration and since $\mathfrak{C}p$ is a weak equivalence, the complex $\mathfrak{C}ITV$ is a cylinder object in the closed model category $\pi\text{-}\mathbf{DM}$. Let $H \in \pi\text{-}\mathbf{DA}(I(TV, d), A)$ be a homotopy between f and g then, $\mathfrak{C}f = \mathfrak{C}H \circ \mathfrak{C}\partial_0$, $\mathfrak{C}g = \mathfrak{C}H \circ \mathfrak{C}\partial_1$ and thus $\mathfrak{C}f \simeq_{\pi} \mathfrak{C}g$.

Appendix B - Equivariant acyclic model theorem.

The proof of Theorem B relies heavily on the π -equivariant acyclic model theorem for cochain functors. We state and prove this theorem.

B.1 Let R be a (ungraded) commutative algebra over the field $\mathbb{F}_p$ and let $\mathbf{C}$ be a category with models $\mathcal{M}$. Consider a contravariant functor $F : \mathbf{C} \rightarrow \mathbf{Coch}_R, A \mapsto FA = \{F^i A\}_{i \in \mathbb{Z}}$ with values in the category of R -cochain complexes (R acting on the left). See [3], [12] or [2] for the definitions of: F admits a unit, F is acyclic on the models and F is corepresentable for the models $\mathcal{M}$. The singular cochains functor $C^n : \mathbf{Top} \rightarrow \mathbf{Mod}_R$ is corepresentable on the models $\{\Delta^k\}_{k \geq 0}$. Observing that a retract of corepresentable functor is corepresentable, we deduce that the functor $X \mapsto N^*X$ is corepresentable on the models $\{\Delta^k\}_{k \geq 0}$. For further use it is interesting to remark here that F is corepresentable on a family of models and if W is an R -free graded module of finite type then so is the functor $A \mapsto \mathrm{Hom}(W, FA)$. Indeed, in this case each $\mathrm{Hom}(W_n, FA)$ is a finite sum on

copies of FA and one concludes using the fact that $\widehat{\prod_{\alpha} F_{\alpha}} = \prod_{\alpha} \hat{F}_{\alpha}$ (see [2]-II for more details).

B.2 We embed the category $\mathbf{Coch}_R$ into the category $\mathbf{Mod}$ of graded (without differential) $\mathbb{F}_p$ -modules. Then, considering the two contravariant functors $F, G : \mathbf{C} \rightarrow \mathbf{Coch}_R$ as functors with values in $\mathbf{Mod}_R$, we denote by $\text{Hom}^i(F, G)$, $i \in \mathbb{Z}$ the sets of natural transformations of degree i . A differential $D : \text{Hom}_R^i(F, G) \rightarrow \text{Hom}_R^{i+1}(F, G)$ is defined by: $(DT)_A = d_{G(A)}T_A - (-1)^i T_A d_{F(A)}$. For our purpose $R = \mathbb{F}_p[\pi] = \mathbb{F}_p[\tau]/(\tau^p - 1)$ then, for any object A of $\mathbf{C}$, $\text{Hom}^i(F, G)$ is a π -module by the rule $\sigma \in \pi, T \in \text{Hom}_R^i(F, G), A \in \mathbf{C}, x \in FA, (\sigma T)_A(x) = \sigma T_A(\sigma^{-1}x)$. The $\mathbb{F}_p$ -module $\text{Hom}_{\pi}(F, G)$ of π -linear transformations is the fixed point set of $\text{Hom}(F, G)$ under this action. The next theorem has been proved in [3].

B.3 Theorem. *Let $\mathbf{C}$ be a category with models $\mathcal{M}_R$ and $F, G : \mathbf{C} \rightarrow \mathbf{Coch}_R$ two contravariant functors with units such that for any $A \in \mathbf{C}$, $F^i A = 0 = G^i A$ if $i < 0$, F is acyclic on the models and G is corepresentable on the models. Then $H^0(\text{Hom}_{\pi}(F, G)) = R$ and $H^i(\text{Hom}_{\pi}(F, G)) = 0$ if $i \neq 0$.*

B.4 Corollary. *If $F : \mathbf{Top} \rightarrow \mathbf{Coch}_{\pi}$ is any functor such that:*

a) $\eta : \mathbb{F}_p \rightarrow F$ is a unit,

b) for any model $\Delta^n, n \geq 0$ there exists a map $\epsilon_n : F(\Delta^n) \rightarrow \mathbb{F}_p$ such that $\epsilon_n \circ \eta_n = id_{\mathbb{F}_p}$

and $\eta_n \circ \epsilon_n \simeq id_{F(\Delta^n)}$ then, $H^i(\text{Hom}_{\pi}(F, \text{Hom}(W, N^*))) = \begin{cases} \mathbb{F}_p[\pi] & i = 0 \\ 0 & i > 0 \end{cases}$.

Proof Recall the (right) π -free acyclic complex W with left π -action defined by $\sigma w = w\sigma^{-1}$, $\sigma \in \pi$ and $w \in W$ and the canonical isomorphism $\text{Hom}_{\pi}(A, \text{Hom}(W, B)) \cong \text{Hom}_{\pi}(W \otimes A, B)$, where (B, d_B) is a (left) π -complex. We precise that π acts trivially on N^*X for any space X . The functor $G = \text{Hom}(W, N^*) : \mathbf{Top} \rightarrow \mathbf{Coch}_{\mathbb{F}_p[\pi]}$, admits a unit η defined as follows: for any space X $\eta_X(\sigma) = \sigma \circ e_0^*$, where $e_0^* \in \text{Hom}(W, N^*X)$ denotes the dual of $e_0 \in W_0$. As remarked in B.1, $G = \text{Hom}(W, N^*)$ is corepresentable on the models Δ^n .

The natural transformation η induces a natural transformation, also denoted $\eta : R = \mathbb{F}_p[\pi] \rightarrow F \otimes R$. Moreover, for any $n \geq 0$ there exists $\epsilon_n = F(\Delta^n) \otimes R \rightarrow R$ such that $\eta_{\Delta^n} \circ \epsilon_{\Delta^n} = id_{\Delta^n}$ and $\epsilon_{\Delta^n} \circ \eta_{\Delta^n} \simeq id_{\Delta^n}$. Since π acts diagonally on $F \otimes R$, the map $\varphi \mapsto (\hat{\varphi} : x \otimes y \mapsto \varphi(x)y)$ is an isomorphism of chain complexes $\text{Hom}_{\pi}(F, G) \cong \text{Hom}_{\pi}(F \otimes R, G)$, for any functor $G : \mathbf{Top} \rightarrow \mathbf{Coch}_{\pi}$. We apply theorem B.3 to end the proof.

References

- [1] D.J. Benson, *Representations and Cohomology II*, Cambridge studies in advanced mathematics **31** - Cambridge University Press - 1991.
- [2] N. Bitjong and J-C. Thomas, *On the cohomology algebra of free loop spaces*, Topology (2001) to appear.

- [3] A.K. Bousfield and V.K.A.M. Gugenheim, *On PL de Rham theory and rational homotopy type*, Memoirs of the Amer. Math. Soc. **179** 1976.
- [4] H. Cartan and S. Eilenberg, *Homological algebra*, Princeton mathematical series **19** - Princeton University Press - 1956.
- [5] S. Eilenberg, *Topological methods in abstract algebra*, Bull. Amer. Math. Soc. **55** (1949) 3-27.
- [6] Y. Félix, S. Halperin and J-C. Thomas, *Adams' cobar equivalence*, Trans. of the Amer. Math. Soc. **329** (1992) 531-549.
- [7] Y. Félix, S. Halperin and J-C. Thomas, *Differential graded algebras in topology*, Handbook of Algebraic Topology, I. James, editor, Chapter 16 - North-Holland - 1995.
- [8] S. Halperin and J. M. Lemaire, *Notions of category in differential algebras*, Algebraic Topology: Rational Homotopy, Springer Lecture Notes in Math. **1318** (1988) 138-153.
- [9] S. Halperin and M. Vigu-Poirrier, *The homology of a free loop space*, Pacific J. of Math. **147** (1991) 311-324.
- [10] J. D.S. Jones, *Cyclic homology and equivariant homology*, Invent. Math. **87** (1987) 403-423.
- [11] J-L. Loday, *Cyclic homology*, Grundlehren der mathematischen Wissenschaften **301** - Springer-Verlag - 1991.
- [12] M. Majewski, *Rational homotopical models and uniqueness*, Memoirs of the Amer. Math. Soc. **143/682**, 2000.
- [13] J.P. May, *A general algebraic approach to Steenrod operations*, Springer Lecture Notes in Math **168** (1970) 153-231.
- [14] S. Mac Lane, *Homology*, Grundlehren der mathematischen Wissenschaften **114** Springer-Verlag - 1963.
- [15] H. J. Munkholm, *The Eilenberg-Moore spectral sequence and strongly homotopy multiplicative maps*, J. of Pure and Appl. Alg. **5** (1974) 1-50.